

Fourier Transform Methods for Option Pricing*

Deng Ding

*Department of Mathematics, University of Macau, Macao,
China*

1. Introduction

Since Bachelier, a French mathematician, first tried to give a mathematical definition for Brownian motion and used it to model the dynamics of stock process in 1900, financial mathematics has developed a lot. Black & Scholes (1973) and Merton (1973) respectively used the geometric Brownian motion (GBM) to model the underlying asset's price process so that they opened the gate of easy ways to compute option prices, which led one of the major breakthroughs of modern finance. Many good ideas have been proposed to model the stock pricing processes since then. Merton (1976) first introduced the jumps into the asset price processes in his seminar paper. More recently, a lot of exponential Lévy models, including Kou's model, Variance Gamma (VG) model, Inverse Gaussian (IG) model, Normal Inverse Gaussian (NIG) model, and CGMY model, etc., were proposed to add jumps in the financial models so that they can describe the statistical properties of financial time series better [e.g. see Cont & Tankov (2003) and references there in]. Also, several stochastic volatility models were presented [e.g. see Heston (1993), Bates (1996), and Duffie *et al* (2000), etc.]. Empirical financial data indicate that these models are usually more consistent with financial markets.

Under assuming that the price of an underlying asset follows a GBM, Black & Scholes (1973) showed that the value of a European option satisfies a boundary problem of heat equation (Black-Scholes equation) so that they derived an explicit formula (Black-Scholes formula) for the value. However, this nice analytic tractability in option pricing can not be carried over to the most exponential Lévy models or the stochastic volatility models for asset returns. Thus, many new approaches, including efficient numerical methods, for option pricing have been proposed. These approaches and methods can be classified into four major groups: The partial integro-differential equation (PIDE) and various numerical methods for such equations; The Monte Carlo methods via the stochastic simulation techniques for underlying asset price processes; Directly numerical integration and various numerical methods via differential integral transforms; Backward stochastic differential equation and its numerical methods. Each of them has its advantages and disadvantages for different financial models and specific applications.

Since Stein & Stein (1991) first used Fourier inversion method to find the distribution of the underlying asset in a stochastic volatility model, the Fourier transform methods have become a very active field of financial mathematics. Heston (1993) applied the characteristic function

*The work was supported by the research grant MYRG136(Y1-L2)-FST11-DD from the Research Committee of University of Macau.

approach to obtain an analytic representation for the valuation of European options in the Fourier domain. Duffie *et al* (2000) offered a comprehensive survey that Fourier transform methods are applicable to a wide range of stochastic processes, the class of exponential affine diffusions [e.g. see Kwok *et al* (2010) and Schmelzle (2010)].

Carr & Madan (1999) pioneered the use of the fast Fourier transform (FFT) technique by mapping the Fourier transform directly to option prices via the characteristic function. Since then, many efficient numerical methods by using FFT techniques have been proposed, and many authors have discussed these methods in rigorous detail. Lee (2004) extended their method significantly and proved an error analysis for these FFT methods. Lewis (2001) generalized this approach to several general payoff functions via the convolution of generalized Fourier transforms.

Recently, some new ideas to improve the Fourier transform methods have been raised. Fang & Oosterlee (2008) proposed the COS method which is based on the Fourier and Fourier-cosine expansion. Feng & Linetsky (2008) presented a new method which involves the relation between Fourier transform and the Hilbert transform, and the Sinc expansion in Hardy spaces. Lord *et al* (2008) extended the FFT-based methods to the CONV method, which is based on the quadrature technique and relies heavily on Fourier transform.

In this chapter, we demonstrate the application of Fourier transform as a very effective tool in option pricing theory. Together with the fast Fourier transform technique and other important properties of Fourier transform, we survey several different methods for pricing European options and some path dependent options under different financial models.

This chapter is organized as follows. In the next section, the mathematical formulations that will be used in this chapter are reviewed. They include a brief discussion on Fourier transform with its important properties; an introduction of discrete Fourier transform and the FFT idea; a definition of characteristic functions; and a brief review on the exponential Lévy models and stochastic volatility models in financial mathematics. In Section 3, several Fourier transform methods for pricing European options are introduced. They include the Black-Scholes type formulas, the FFT methods for signal underlying asset and for multi underlying assets, and the Fourier expansion methods. In Section 4, three new Fourier transform methods for pricing path dependent options are considered. They are the CONV method for pricing Bermudan barrier option, the COS method for pricing Bermudan barrier option, and the fast Hilbert transform method for pricing barrier option.

2. Fourier transforms and characteristic functions

2.1 Fourier transforms

Let $g(x)$ be a piecewise continuous real function over \mathbb{R} which satisfies the integrability condition:

$$\int_{-\infty}^{\infty} |g(x)| dx < \infty.$$

The *Fourier transform* of $g(x)$ is defined by

$$\mathcal{F}g(u) = \int_{-\infty}^{\infty} e^{iux} g(x) dx, \quad u \in \mathbb{R}, \quad (1)$$

where $i = \sqrt{-1}$ is the imaginary unit. Given the function $\mathcal{F}g(u)$, the function $g(x)$ can be recovered by the *Fourier inversion formula*:

$$g(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-iux} \mathcal{F}g(u) du, \quad x \in \mathbb{R}. \tag{2}$$

Sometime it may be more convenient to consider the *generalized Fourier transform* on the complex plane. Let $a, b \in \mathbb{R}$ with $a < b$. Assume

$$\int_{-\infty}^{\infty} e^{-ax} |g(x)| dx < \infty$$

and

$$\int_{-\infty}^{\infty} e^{-bx} |g(x)| dx < \infty.$$

Then, the generalized Fourier transform:

$$\mathcal{F}g(z) = \int_{-\infty}^{\infty} e^{izx} g(x) dx, \quad z \in \mathbb{C},$$

exists and is analytic for all z in the strip $\mathbb{S} = \{z \in \mathbb{C} : a < \text{Im}\{z\} < b\}$. Moreover, within this strip the generalized Fourier transform may be inverted by integrating along a straight line paralleled to the real axis:

$$g(x) = \frac{1}{2\pi} \int_{iw-\infty}^{iw+\infty} e^{-ixz} \mathcal{F}g(z) dz, \quad x \in \mathbb{R}, \tag{3}$$

with $a < w < b$. Here, $\text{Im}\{\cdot\}$ denotes taking the imaginary part of argument.

The following properties of Fourier transform are useful in this chapter.

P1. Differentiation: $\mathcal{F}(g'(x))(u) = -iu\mathcal{F}g(u)$, where $g'(x)$ is the derivative of $g(x)$.

P2. Modulation: $\mathcal{F}(e^{\lambda x} g(x))(u) = \mathcal{F}g(u - i\lambda)$, where λ is a constant.

P3. Convolution: $\mathcal{F}(f * g)(u) = \mathcal{F}f(u)\mathcal{F}g(u)$, where $f * g$ is the convolution of $f(x)$ and $g(x)$, which is defined by

$$(f * g)(x) = \int_{-\infty}^{\infty} f(y)g(x - y) dy.$$

P4. Relation with Hilbert transform: $\mathcal{F}(\text{sgn} \cdot g)(u) = i\mathcal{H}(\mathcal{F}g)(u)$, where $\text{sgn}(x)$ is the signum function, and \mathcal{H} is the Hilbert transform, which is defined by the Cauchy principal value integral:

$$\mathcal{H}f(u) = \frac{1}{\pi} P.V. \int_{-\infty}^{\infty} \frac{f(x)}{u - x} dx.$$

P5. Parseval relation: $\langle f, g \rangle = \frac{1}{2\pi} \langle \mathcal{F}f(u), \mathcal{F}g(u) \rangle$, where $\langle \cdot, \cdot \rangle$ is the inner product of two square-integrable function defined by

$$\langle f, g \rangle = \int_{-\infty}^{\infty} f(x)g(x) dx.$$

2.2 Fast Fourier transforms

Let $N = 2^L$ be a power of 2, and let $x = (x_0, x_1, \dots, x_{N-1})^T$ be a given N -dimensional vector, where v^T presents the transpose of a vector v . The *discrete Fourier transform* of x is another N -dimensional vector $\mathcal{D}x = (y_0, y_1, \dots, y_{N-1})^T$ is defined by

$$y_p = \sum_{j=0}^{N-1} e^{i\frac{2\pi}{N}jp} x_j, \quad p = 0, 1, \dots, N-1. \tag{4}$$

Denote A_N an $N \times N$ matrix whose $(p, j)^{\text{th}}$ entry is given by

$$a_{pj} = e^{i\frac{2\pi}{N}jp}, \quad j, p = 0, 1, \dots, N-1.$$

Then, the *discrete Fourier transform* of x is given by

$$\mathcal{D}x = A_N x.$$

It is clear that the computation to find $\mathcal{D}x$ requires N^2 steps. However, the *fast Fourier technique* (FFT) would require only $\frac{1}{2}NL = \frac{1}{2}N \log_2 N$ steps. The idea behind the FFT algorithm is to take advantage of the periodicity property of N roots of unity. For the given N -dimensional vector x , denote $x' = (x_0, x_2, \dots, x_{N-2})^T$ and $x'' = (x_1, x_3, \dots, x_{N-1})^T$ two $N/2$ -dimensional vectors. Then, we find

$$y' = A_M x' \quad \text{and} \quad y'' = A_M x'',$$

where $M = N/2$, $y' = (y'_0, y'_1, \dots, y'_{M-1})^T$ and $y'' = (y''_0, y''_1, \dots, y''_{M-1})^T$, and the matrix $A_M = [a_{jk}]_{M \times M}$ given by

$$a_{pj} = e^{i\frac{2\pi}{N}jp}, \quad j, p = 0, 1, \dots, M-1.$$

It is easy to verify that the vector $\mathcal{D}x$, which is defined in (4), now is given by

$$\begin{aligned} y_p &= y'_p + e^{i\frac{2\pi}{N}p} y''_p, \quad p = 0, 1, \dots, M-1, \\ y_{M+p} &= y'_p - e^{i\frac{2\pi}{N}p} y''_p, \quad p = 0, 1, \dots, M-1. \end{aligned}$$

Instead of performing the matrix-vector multiplication $A_N x$, we now only need to perform two matrix-vector multiplications $A_M x'$ and $A_M x''$ so that the number of operations is reduced from N^2 to $2(N/2)^2 = N^2/2$. The same procedure of reducing the length of the vector by half can be applied repeatedly. Using this FFT algorithm, the total number of operations is reduced from $O(N^2)$ to $O(N \log_2 N)$. The similar FFT algorithm also can be used to calculate the *discrete Fourier inversion transform*:

$$x_p = \sum_{j=0}^{N-1} e^{-i\frac{2\pi}{N}jp} y_j, \quad p = 0, 1, \dots, N-1.$$

2.3 Characteristic functions

Let X be a random variable having the density function $f(x)$. Then, the *characteristic function* of X is defined by

$$\varphi_X(u) = \mathbb{E}[e^{iuX}] = \int_{-\infty}^{\infty} e^{iux} f(x) dx, \quad u \in \mathbb{R},$$

i.e. it is the Fourier transform of the density $f(X)$. Here $\mathbb{E}[\cdot]$ denotes the mathematical expectation. The characteristic functions are uniformly continuous and non-negative definite. Also they possess the following properties:

P6.If X and Y are independent, then $\varphi_{X+Y}(u) = \varphi_X(u)\varphi_Y(u)$.

P7.If $a, b \in \mathbb{R}$ and $Y = aX + b$, then $\varphi_Y(u) = e^{ibu}\varphi_X(au)$.

P8. X and Y have the same distribution function if and only if they have the same characteristic function.

2.4 Exponential Lévy models

An adapted stochastic process X_t with $X_0 = 0$ is called a *Lévy process* if it has independent and stationary increments, and it is continuous in probability. Moreover, every Lévy process has a right continuous with left limits (càdlàg) version which is also a Lévy process. We always work with this càdlàg version.

Let X_t be a Lévy process, and let \mathcal{B}_0 be the set family of all Borel sets $U \subset \mathbb{R}$ whose closure \bar{U} does not contain 0. Set

$$\mu((0, t], U) = \sum_{0 < s \leq t} 1_U(\Delta X_s), \quad t > 0, U \in \mathcal{B}_0,$$

and denote $\mu(\{0\}, U) \equiv 0$. Then $\mu(dt, dx)$ is a σ -finite random measure on $\mathcal{B}(\mathbb{R}_+) \times \mathcal{B}_0$, which is called the *jump measure* of X_t . Set $\nu(U) = \mathbb{E}[\mu([0, 1], U)]$ for any $U \in \mathcal{B}_0$. It is clear that $\nu(dx)$ is a σ -finite measure on \mathcal{B}_0 , which is called the *Lévy measure* of X_t . Then, the characteristic function of Lévy process X_t has the *Lévy-Khintchine representation*:

$$\varphi_t(u) = e^{t\psi(u)}, \quad u \in \mathbb{R}, \tag{5}$$

where the characteristic exponent function:

$$\psi(u) = i\alpha u - \frac{1}{2}\sigma^2 u^2 + \int_{\mathbb{R}} (e^{iux} - 1 - iux1_{\{|x| \leq 1\}}(x))\nu(dx). \tag{6}$$

Here the triple $(\alpha, \sigma^2, \nu(dx))$ is called the Lévy triple of X_t . The proof of this Lévy-Khintchine decomposition is based on the famous *Lévy-Itô decomposition*, and the detail can be found in Cont & Tankov (2003). Also a Lévy process is a strong Markov process, and a semimartingale. Furthermore, under the condition: $\int_{\{|x| \geq 1\}} e^x \nu(dx) < \infty$, the exponential e^{X_t} is a martingale if and only if

$$\alpha + \frac{1}{2}\sigma^2 + \int_{\mathbb{R}} (e^x - 1 - x1_{\{|x| \leq 1\}}(x))\nu(dx) = 0.$$

To ensure positivity as well as the independence and stationarity of log-returns, in financial mathematics, the price process of the underlying asset is usually modeled as the exponential of a Lévy process X_t :

$$S_t = S_0 e^{X_t}, \quad t \geq 0. \tag{7}$$

There are many models in the financing modeling literature simply correspond to different choices for σ^2 and $\nu(dx)$ to the Lévy process X_t .

The cases that $\sigma^2 \neq 0$ and $\nu(dx)$ is a finite measure, i.e. X_t is a Lévy jump diffusion:

$$X_t = \mu t + \sigma B_t + \sum_{k=1}^{N_t} \tilde{\zeta}_k, \quad t \geq 0,$$

where B_t is a standard Brownian motion, $\{\tilde{\zeta}_k\}$ is a sequence of independent and identically distributed random variables with common density $f_{\tilde{\zeta}}(x)$, and N_t is a Poisson process of intensity λ , such that B_t , $\{\tilde{\zeta}_k\}$ and N_t are mutually independent. In these models, the Lévy triple is given by

$$\alpha = \mu + \lambda \int_{\{|x|<1\}} x f_{\tilde{\zeta}}(x) dx, \quad \sigma^2 \quad \text{and} \quad \nu(dx) = \lambda f_{\tilde{\zeta}}(x) dx.$$

These models explain the jump part as the market responses to outside news: good news and bad news. The news arrives according to the Poisson process N_t , and the price changes in response according to the jump size $\tilde{\zeta}_k$.

- The earliest of these models is due to Merton (1976). In this model, $\tilde{\zeta}_{kS}$ are normally distributed, with mean μ_j and standard deviation σ_j , so that the characteristic function of X_t is given by

$$\varphi_t(u) = \exp \left\{ i\alpha t u - \frac{1}{2} \sigma^2 t u^2 + \lambda t \left(e^{i\mu_j u - \frac{1}{2} \sigma_j^2 u^2} - 1 \right) \right\}.$$

- The second one belongs to Kou (2002). In this model, $\tilde{\zeta}_{kS}$ are non-symmetric double exponentially distributed, and the characteristic function of X_t is given by

$$\varphi_t(u) = \exp \left\{ i\alpha t u - \frac{1}{2} \sigma^2 t u^2 + \lambda t \left(\frac{1 - \eta^2}{1 + \eta^2 u^2} e^{i\kappa u} - 1 \right) \right\},$$

where η and κ are parameters.

The cases that $\sigma^2 = 0$ and $\nu(\mathbb{R} \setminus \{0\}) = \infty$, i.e., the models have infinite activity in jumps. We only list some examples here.

- The VG model, which is proposed by Madan *et al* (1998): X_t is a variance gamma process with parameters σ, κ and θ , and its characteristic function is given by:

$$\varphi_t(u) = \left(1 - i\theta\kappa u + \frac{1}{2} \sigma^2 \kappa u^2 \right)^{-t/\kappa}.$$

- The NIG model, which is presented by Barndorff-Nielsen (1997): X_t is a normal inverse Gaussian process with parameters α, β and δ , and its characteristic function is given by:

$$\varphi_t(u) = \exp \left\{ -\delta t \left(\sqrt{\alpha^2 - (\beta + iu)^2} - \sqrt{\alpha^2 - \beta^2} \right) \right\}.$$

- The CGMY model, which is defined by Carr *et al* (2002): X_t is a CGMY process whose characteristic function is given by

$$\varphi_t(u) = \exp \left\{ t \text{CG}(-Y) \left[(M - iu)^Y - M^Y + (G + iu)^Y - G^Y \right] \right\},$$

where the parameters C, G and M are nonnegative, $Y < 2$, and $\Gamma(x)$ is the gamma function.

- The Finite Moment Log Stable model, which is considered by Carr & Wu (2003): X_t is a finite moment log stable process whose characteristic function is given by

$$\varphi_t(u) = \exp \left\{ i\omega tu - (i\sigma u)^\alpha t \sec \frac{\pi\alpha}{2} \right\},$$

where ω , α and σ are parameters.

2.5 Stochastic volatility models

Although the class of Lévy processes is quite rich, it is sometime insufficient for multi-period financial modeling. Several models combining jumps and stochastic volatility have been appeared in the literature. We only list two such models here.

- The Heston’s stochastic volatility model [Heston (1993)]: In this model, the price process S_t is defined by the system of stochastic differential equations (SDEs):

$$\begin{cases} dS_t = \mu S_t dt + \sqrt{V_t} S_t dB_t^S, \\ dV_t = \kappa(\theta - V_t)dt + \sigma\sqrt{V_t} dB_t^V, \end{cases}$$

with the initial values $S_0 = s_0$ and $V_0 = v_0$, where B_t^S and B_t^V are two standard Brownian motions with correlation ρ (i.e. $\langle B^S, B^V \rangle_t = \rho t$), κ is the mean reversion, θ is the long-run variance level, and σ is the volatility-of-volatility parameter. Although S_t is no longer a Lévy process, under the risk neutral probability the characteristic function of the log-price process $X_t = \log S_t$ is known in closed form:

$$\varphi_t(u) = \frac{\exp \left\{ \frac{\kappa\theta t(\kappa - i\rho\sigma u)}{\sigma^2} + irtu + ix_0u \right\}}{\left(\cosh \frac{\gamma t}{2} + \frac{\kappa - i\rho\sigma u}{\gamma} \sinh \frac{\gamma t}{2} \right)^{2\kappa\theta/\sigma^2}} \exp \left\{ - \frac{(u^2 + iu)v_0}{\gamma \coth \frac{\gamma t}{2} + \kappa - i\rho\sigma u} \right\},$$

where $x_0 = \log s_0$, r is the interest rate, and $\gamma = \sqrt{\sigma^2(u^2 + iu) + (\kappa - i\rho\sigma u)^2}$.

- The Baste’s stochastic volatility model [Bates (1996)]: In this model, the price process S_t and the variance process V_t are given by the system of SDEs:

$$\begin{cases} dS_t = \mu S_{t-} dt + \sqrt{V_t} S_{t-} dB_t^S + S_{t-} dZ_t, \\ dV_t = \kappa(\theta - V_t)dt + \sigma\sqrt{V_t} dB_t^V, \end{cases}$$

where B_t^S and B_t^V are two standard Brownian motions with correlation ρ , and Z_t is a compound Poisson process with intensity λ and log-normal distribution of jump sizes such that if k is its jump size then

$$\log(1 + k) \sim N \left(\log(1 + \bar{k}) - \frac{1}{2}\delta^2, \delta^2 \right).$$

Similarly to the Heston’s model, the characteristic function of $X_t = \log S_t$ is known in closed form:

$$\varphi_t(u) = \varphi_t^D(u) \cdot \varphi_t^J(u),$$

where the characteristic function of the diffusion part:

$$\varphi_t^D(u) = \frac{\exp \left\{ \frac{\kappa\theta t(\kappa - i\rho\sigma u)}{\sigma^2} + i(r - \lambda\bar{k})tu + ix_0u \right\}}{\left(\cosh \frac{\gamma t}{2} + \frac{\kappa - i\rho\sigma u}{\gamma} \sinh \frac{\gamma t}{2} \right)^{2\kappa\theta/\sigma^2}} \exp \left\{ - \frac{(u^2 + iu)v_0}{\gamma \coth \frac{\gamma t}{2} + \kappa - i\rho\sigma u} \right\},$$

and the characteristic function of the jump part:

$$\varphi_t^J(u) = \exp \left\{ t\lambda \left(e^{-\frac{1}{2}\delta^2 u^2 + i(\log(1+\bar{k}) - \frac{1}{2}\delta^2)u} - 1 \right) \right\}.$$

3. Fourier transform methods for pricing European options

3.1 Black-Scholes type formulas

Gurldand (1948) used the Convolution property P3 to derive an formula for calculation of the distribution function $F(x)$ from the characteristic function $\varphi(u)$:

$$F(x) = \frac{1}{2} + \frac{1}{2\pi} \int_0^\infty \frac{e^{iux} \varphi(-u) - e^{-iux} \varphi(u)}{iu} du.$$

Shephard (1991) gave a simple proof of this formula. From this formula by a simple calculation we can get an important formula for Fourier inversion method:

$$\mathbb{P}(X > x) = 1 - F(x) = \frac{1}{2} + \frac{1}{\pi} \int_0^\infty \operatorname{Re} \left(\frac{e^{-iux} \varphi(u)}{iu} \right) du. \tag{8}$$

Here $\operatorname{Re}\{\cdot\}$ denotes taking the real part of argument. Inside the field of finance, this Fourier inversion method was first considered by Stein & Stein (1991) to find the distribution of the underlying asset price in a stochastic volatility model. Heston (1993) applied the formula (8), to obtain a Black-Scholes type formula for pricing a European call option in the stochastic volatility model.

In fact, with help of the risk-neutral valuation, the price of a European call with spot price S_0 and strike price K is given by

$$\begin{aligned} V(S_0, T; K) &= e^{-rT} \mathbb{E}[(S_T - K)^+] = e^{-rT} \int_{-\infty}^\infty (e^x - e^k)^+ f_T(x) dx \\ &= \int_k^\infty e^{-rT+x} f_T(x) dx - e^{-rT} K \int_k^\infty f_T(x) dx = I_1 - e^{-rT} K I_2, \end{aligned}$$

where r is the interest rate, T is the maturity, $k = \log K$, and $f_T(x)$ is the density function of $X_T = \log S_T$. It is clear that the second integral is the probability: $\Pi_2 = \mathbb{P}(X_t > k)$. By the Fourier inversion formula (8) we have

$$\Pi_2 = \frac{1}{2} + \frac{1}{\pi} \int_0^\infty \operatorname{Re} \left(\frac{e^{-iuk} \varphi_T(u)}{iu} \right) du,$$

where $\varphi_T(u)$ is the characteristic function of X_T . By a change of probability measure, we can verify that the second integral is give by $I_1 = S_0 \Pi_1$, where

$$\Pi_1 = \frac{1}{2} + \frac{1}{\pi} \int_0^\infty \operatorname{Re} \left(\frac{e^{-iuk} \varphi_T(u - i)}{iu \varphi_T(-i)} \right) du,$$

Thus, the price of the European call now is given by a Black-Scholes type formula:

$$V(S_0, T; K) = S_0 \Pi_1 - K e^{-r(T-t)} \Pi_2. \tag{9}$$

Beginning with Heston’s work, many authors used the Fourier inversion methods to solve advanced valuation problems [e.g. see Bates (1996), Schmelzle (2010), and references therein]. As mentioned by Carr & Madan (1999), the numerical integrals based on these methods are generally much faster than finite difference solutions to PDEs or PIDEs. Unfortunately, the FFT cannot be used to evaluate these integrals, since the integrands are singular at the required evaluation point $u = 0$.

Recently, several authors applied the Property P5 (Parseval relation) to derive the Fourier inversion formulas in option pricing [e.g. see Dufresne *et al* (2009)]. We have well known that, under the risk-neutral probability, the option price $V(S_0, T)$ with the terminal payoff $H(S_T)$ is given by

$$V(S_0, T) = e^{-rT} \mathbb{E}[H(S_T)] = e^{-rT} \int_{-\infty}^{\infty} H(x) p_T(x) dx = e^{-rT} \langle H(x), p_T(x) \rangle,$$

where $p_T(x)$ is the density function of S_T . By the Parseval relation we have

$$V(S_0, T) = \frac{e^{-rT}}{2\pi} \langle \mathcal{F}H(u), \mathcal{F}p_T(u) \rangle.$$

3.2 Fast Fourier transform methods - single asset

1. The method of Carr and Madan. Carr & Madan (1999) developed a different method designed to use the fast Fourier transform to price options. They introduced a new technique to calculate the Fourier transform of a modified call option price with respect to the logarithmic strike price so that the fast Fourier transform can be applied to calculate the integrals.

Consider a European call with the maturity T and the strike price K , which is written on a stock whose price process is $S_t = e^{rt+X_t}$, under a risk-neutral probability. Let $f_T(x)$ be the density of X_T . Consider the price of a call option:

$$V_T(k) = e^{-rT} \mathbb{E}[(e^{X_T} - e^k)^+] = e^{-rT} \int_k^{\infty} (e^x - e^k) f_T(x) dx, \tag{10}$$

where $k = \ln K$ is the log strike price. Note that $V_T(k) \rightarrow S_0 = 1$ as $k \rightarrow -\infty$, and the function $V_T(k)$ is not square-integrable. Thus, we cannot express the Fourier transform in strike in terms of the characteristic function $\varphi_T(u)$ of X_T and then find a range of strikes by Fourier inversion.

To obtain a square-integrable function we consider the modified call price:

$$\tilde{V}_T(k : \alpha) = e^{\alpha k} V_T(k)$$

for some $\alpha > 0$, which is chosen to improve the integrability. Carr & Madan (1999) showed that a sufficient condition for square-integrability of $\tilde{V}(k)$ is given by

$$\int_{-\infty}^{\infty} f_T(x) e^{(1+\alpha)x} dx < \infty.$$

Consider the Fourier transform of $\tilde{V}(k : \alpha)$:

$$\psi_T(v : \alpha) = \mathcal{F}\tilde{V}_T(k : \alpha)(v) = \int_{-\infty}^{\infty} e^{ivk} \tilde{V}_T(k : \alpha) dk.$$

By (10) and the definition of characteristic functions we have

$$\begin{aligned} \psi_T(v : \alpha) &= e^{-rT} \int_{-\infty}^{\infty} f_T(x) dx \int_{-\infty}^x (e^{x+\alpha k} - e^{(1+\alpha)k}) e^{ivk} dk \\ &= \frac{e^{-rT} \varphi_T(v - i(\alpha + 1))}{\alpha^2 + \alpha - v^2 + i(2\alpha + 1)v'} \end{aligned} \tag{11}$$

where $\varphi_T(u)$ is the characteristic function of X_T . Using the Fourier inverse transform we get

$$V_T(k) = e^{-\alpha k} \tilde{V}(k : \alpha) = \frac{e^{-\alpha k}}{2\pi} \int_{-\infty}^{\infty} e^{-ivk} \psi_T(v : \alpha) dv.$$

Note that $\psi_T(v : \alpha)$ is odd in its imaginary part and even in its real part. Since $V(k)$ is real we have

$$V_T(k) = \frac{e^{-\alpha k}}{\pi} \int_0^{\infty} e^{-ivk} \psi_T(v : \alpha) dv. \tag{12}$$

Now this integral can be evaluated by the numerical approximation using the trapezoidal rule:

$$V_T(k_p) \approx \frac{e^{-\alpha k_p}}{2\pi} \sum_{j=0}^{N-1} e^{-iv_j k_p} \psi_T(v_j : \alpha) \Delta v, \quad p = 0, 1, \dots, N-1.$$

to apply the FFT algorithm, we set N as a power of 2, and define the grid points:

$$\begin{cases} k_p = -\frac{1}{2}N\Delta k + p\Delta k, & p = 0, 1, \dots, N-1, \\ v_j = -\frac{1}{2}N\Delta v + j\Delta v, & j = 0, 1, \dots, N-1, \end{cases}$$

with the step sizes Δk and Δv , which satisfy the Nyquist relation: $\Delta k \Delta v = 2\pi/N$. Then, the numerical approximation can be expressed as

$$V_T(k_p) \approx \frac{e^{-\alpha k_p}}{2\pi} \sum_{j=0}^{N-1} (-1)^{j+p} e^{-i\frac{2\pi}{N}jp} \psi_T(v_j : \alpha) \Delta v, \quad p = 0, 1, \dots, N-1,$$

which can be efficiently computed by using FFT algorithm.

Many authors have discussed this Carr and Madan’s pricing method in rigorous detail. Lee (2004) extended this method significantly and proved an error analysis for this FFT method. On the other hand, numerical experiments show that this FFT method has a quite large error when the strike price K is small, and its stability is very dependent on the choice of the damping exponential factor α . Ding & U (2010) presented a modified FFT-based method to overcome these disadvantages.

2. The method of Lewis. Lewis (2001) introduced an option pricing method which generalizes previous work on Fourier transform methods. The main idea of his method is to express the option price via the convolution of generalized Fourier transforms, and then, to apply the property P5 (Parseval relation) to obtain the generalized Fourier transform of option price.

Consider a European type option whose payoff is $H(S_T)$ at maturity T , where $S_t = S_0 e^{rt+X_t}$ is the stock price process under the risk neutral probability. To proceed, we assume that X_t is a Lévy process having the analytic characteristic function $\varphi_T(z)$, which is regular in the strip

$S_X = \{z : a < \text{Im}\{z\} < b\}$, and $h(x) = H(e^{x+rT})$ is Fourier integrable in some strip S_h such that

$$S_V = \bar{S}_X \cap S_h \neq \emptyset,$$

where \bar{S}_X is the complex conjugate set of S_X .

Let $s = \log S_0$ denote the logarithm of current stock value. Then, the option price is given by

$$V(s) = e^{-rT} \mathbb{E} \left[H(e^{s+rT+X_T}) \right] = e^{-rT} \int_{-\infty}^{\infty} H(e^{s+rT+x}) f_T(x) dx,$$

where $f_T(x)$ is density function of X_T . Under the assumption we can compute the generalized Fourier transform of $V(s)$ by the Parseval relation:

$$\begin{aligned} \int_{-\infty}^{\infty} e^{izs} V(s) ds &= e^{-rT} \int_{-\infty}^{\infty} f_T(x) dx \int_{-\infty}^{\infty} e^{izs} H(e^{s+rT+x}) ds \\ &= e^{-rT} \int_{-\infty}^{\infty} e^{-izx} f_T(x) dx \int_{-\infty}^{\infty} e^{izy} H(e^{y+rT}) dy, \end{aligned}$$

for all $z \in S$. Finally we obtain

$$\mathcal{F}V(z) = e^{-rT} \varphi_T(-z) \mathcal{F}h(z), \quad z \in S_V.$$

Option prices can now be given by the generalized Fourier inversion formula (3):

$$V(s) = \frac{e^{-rT}}{2\pi} \int_{iw-\infty}^{iw+\infty} \varphi_T(-z) \mathcal{F}h(z) dz,$$

with $a < w < b$, for a range of initial values s .

For a European call option, the function $h(x) = (e^{rT+x} - e^k)^+$, is Fourier integrable in the region $\text{Im}\{z\} > 1$, where its generalized Fourier transform can be computed explicitly:

$$\mathcal{F}h(z) = \int_{-\infty}^{\infty} e^{ixz} (e^{x+rT} - e^k)^+ dx = \frac{e^{k+iz(k-rT)}}{iz(iz+1)}.$$

Hence, the generalized Fourier transform of call option price takes the form:

$$\mathcal{F}V(z) = \varphi_T(-z) \frac{e^{(1+iz)(k-rT)}}{iz(iz+1)}, \quad 1 < \text{Im}\{z\} < 1 + \alpha.$$

and hence, the option price is given by

$$V(s) = \frac{e^{ws+(1-w)(k-rT)}}{2\pi} \int_{-\infty}^{\infty} \frac{\varphi_T(-z) e^{iu(k-rT-s)}}{(iu-w)(1+iu-w)} du,$$

for some $1 < w < 1 + \alpha$. The integral in this formula can be approximated by using the FFT algorithm as the formula (12). However, the choice of w is a delicate issue because choosing big w leads to slower decay rates at infinity and bigger truncation errors, and while w is close to one the denominator diverges and the discretization error becomes to large (see Chapter 11 in Cont & Tankov (2003)). On the other hand, Lewis (2001) also listed the generalized Fourier transforms $\mathcal{F}h(z)$ and the strip S_h for various claims, for instance, the put option, the covered call or the cash-secured put, etc.

3.3 Fast Fourier transform methods - multi assets

1. The most direct extension of the Carr and Madan method to the multi assets model is to price the correlation option, whose payoff at the maturity T is defined by

$$H(S_{1T}, S_{2T}) = (S_{1T} - K_1)^+(S_{2T} - K_2)^+,$$

where K_1 and K_2 are strike prices [see, e.g. Kwok *et al* (2010)]. Define $X_{it} = \log S_{it}$ and $k_i = \log K_i, i = 1, 2$, and let $f_T(x_1, x_2)$ be the joint density of X_{1T} and X_{2T} under the risk neutral probability. Then, the characteristic function of X_{1T} and X_{2T} is defined by the following two-dimensional Fourier transform:

$$\varphi_T(u_1, u_2) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i(u_1x_1 + u_2x_2)} f_T(x_1, x_2) dx_1 dx_2.$$

Consider the price of the correlation option:

$$V_T(k_1, k_2) = e^{-rT} \int_{k_2}^{\infty} \int_{k_1}^{\infty} (e^{x_1} - e^{k_1})(e^{x_2} - e^{k_2}) f_T(x_1, x_2) dx_1 dx_2.$$

Following the Carr and Madam method we consider the modified call price:

$$\tilde{V}_T(k_1, k_2 : \alpha_1, \alpha_2) = e^{\alpha_1 k_2 + \alpha_2 k_1} V_T(k_1, k_2),$$

for some parameters $\alpha_1 > 0$ and $\alpha_2 > 0$, which are chosen such that this modified price is square integrable for negative value of k_1 and k_2 . Then, by a direct integral the Fourier transform of $\tilde{V}(k_1, k_2)$ is given by

$$\begin{aligned} \psi_T(v_1, v_2) &= \mathcal{F}\tilde{V}_T(k_1, k_2 : \alpha_1, \alpha_2) \\ &= \frac{e^{-rT} \varphi_T(v_1 - i(\alpha_1 + 1), v_2 - i(\alpha_2 + 1))}{(\alpha_1 + iv_1)(\alpha_1 + 1 + iv_1)(\alpha_2 + iv_2)(\alpha_2 + 1 + iv_2)}. \end{aligned}$$

Applying the Fourier inversion on $\psi_T(v_1, v_2)$ and using the numerical approximation of two-dimensional Fourier inversion integral we have

$$\begin{aligned} V_T(k_p^1, k_q^2) &= \frac{e^{-\alpha_1 k_p^1 - \alpha_2 k_q^2}}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-i(v_1 k_p^1 + v_2 k_q^2)} \psi_T(v_1, v_2) dv_1 dv_2 \\ &\approx \frac{e^{-\alpha_1 k_p^1 - \alpha_2 k_q^2}}{(2\pi)^2} \sum_{m=0}^{N-1} \sum_{n=0}^{N-1} e^{-i(v_m^1 k_p^1 + v_n^2 k_q^2)} \psi_T(v_m^1, v_n^2) \Delta v^1 \Delta v^2, \end{aligned}$$

for $p, q = 0, 1, \dots, N-1$. To apply the two-dimensional of the FFT algorithm we set

$$v_m^1 = (m - \frac{N}{2})\Delta v^1, \quad v_n^2 = (n - \frac{N}{2})\Delta v^2, \quad m, n = 0, 1, \dots, N-1,$$

and

$$k_p^1 = (p - \frac{N}{2})\Delta k^1, \quad k_q^2 = (q - \frac{N}{2})\Delta k^2, \quad p, q = 0, 1, \dots, N-1,$$

where N is a power of 2, and the step sizes $\Delta v^1, \Delta v^2, \Delta k^1$, and Δk^2 observe the Nyquist relations: $\Delta v^1 \Delta k^1 = \Delta v^2 \Delta k^2 = 2\pi/N$. Dempster & Hong (2002) shown that the numerical approximation now is given by

$$V_T(k_p^1, k_q^2) \approx \frac{e^{-\alpha_1 k_p^1 - \alpha_2 k_q^2}}{(2\pi)^2} \Psi_T(k_p^1, k_q^2),$$

for $k_p^1 \neq k_q^2$, where

$$\Psi_T(k_p^1, k_q^2) = (-1)^{p+q} \sum_{m=0}^{N-1} \sum_{n=0}^{N-1} e^{-i\frac{2\pi}{N}(mp+nq)} ((-1)^{m+n} \psi_T(v_m^1, v_n^2) \Delta v^1 \Delta v^2),$$

which can be computed via the FFT algorithm.

There are some other types of terminal payoff functions that admit analytic representation of the Fourier transform of the damped option price [see e.g. Eberlein *et al* (2010), Lee (2004), and references therein]. However, to derive the FFT pricing algorithm for the spread option:

$$H(S_{1T}, S_{2T}) = (S_{1T} - S_{2T} - K)^+, \tag{13}$$

Dempster & Hong (2002) approximated the exercise region of $H(S_{1T}, S_{2T})$ by a combination of rectangular strips.

2. Hurd & Zhou (2010) proposed an alternative approach to pricing the European spread option (13) under the three-factor SV model and exponential Lévy models. Let $h(x_1, x_2)$ be the terminal spread option payoff with the strike price $K = 1$, i.e.

$$h(x_1, x_2) = (e^{x_1} - e^{x_2} - 1)^+. \tag{14}$$

And denote $\Gamma(z)$ the complex gamma function defined by

$$\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt, \quad \text{Re}\{z\} > 0.$$

Then, the method of Hurd and Zhou mainly relies on the following Fourier representation of the payoff function $h(x_1, x_2)$:

$$h(x_1, x_2) = \frac{1}{(2\pi)^2} \int_{i\epsilon_2 - \infty}^{i\epsilon_2 + \infty} \int_{i\epsilon_1 - \infty}^{i\epsilon_1 + \infty} e^{i(u_1 x_1 + u_2 x_2)} \hat{h}(u_1, u_2) du_1 du_2, \tag{15}$$

where ϵ_1 and ϵ_2 are any two given real numbers with $\epsilon_2 > 0$ and $\epsilon_1 + \epsilon_2 < -1$, and

$$\hat{h}(u_1, u_2) = \frac{\Gamma(i(u_1 + u_2) - 1) \Gamma(-iu_2)}{\Gamma(iu_1 + 1)}.$$

The proof of this formula can be found in the appendix of Hurd & Zhou (2010). Let $X_{1t} = \log S_{1t}$ and $X_{2t} = \log S_{2t}$ with $X_{10} = x_1$ and $X_{20} = x_2$, and let $\varphi_T(u_1, u_2)$ be the characteristic function of $X_{1T} - x_1$ and $X_{2T} - x_2$. Then, we have

$$\mathbb{E}[e^{i(u_1 X_{1T} + u_2 X_{2T})}] = e^{i(u_1 x_1 + u_2 x_2)} \varphi_T(u_1, u_2).$$

Now, using the formula (16), the price of the spread option (14) is expressed as an explicit two-dimensional Fourier inversion transform:

$$\begin{aligned} V_T(x_1, x_2) &= e^{-rT} \mathbb{E}[(e^{X_{1T}} - e^{X_{2T}} - 1)^+] \\ &= e^{-rT} \mathbb{E}\left[\frac{1}{(2\pi)^2} \int_{i\epsilon_2 - \infty}^{i\epsilon_2 + \infty} \int_{i\epsilon_1 - \infty}^{i\epsilon_1 + \infty} e^{i(u_1 X_{1T} + u_2 X_{2T})} \hat{h}(u_1, u_2) du_1 du_2\right] \\ &= \frac{e^{-rT}}{(2\pi)^2} \int_{i\epsilon_2 - \infty}^{i\epsilon_2 + \infty} \int_{i\epsilon_1 - \infty}^{i\epsilon_1 + \infty} \mathbb{E}[e^{i(u_1 X_{1T} + u_2 X_{2T})}] \hat{h}(u_1, u_2) du_1 du_2 \\ &= \frac{e^{-rT}}{(2\pi)^2} \int_{i\epsilon_2 - \infty}^{i\epsilon_2 + \infty} \int_{i\epsilon_1 - \infty}^{i\epsilon_1 + \infty} e^{i(u_1 x_1 + u_2 x_2)} \varphi_T(u_1, u_2) \hat{h}(u_1, u_2) du_1 du_2. \end{aligned}$$

This two-dimensional Fourier inversion integral can be numerically computed by the usual FFT calculations. Since this approach does not require an approximation of the exercise region, it is considered to be more computationally efficient.

3.4 Fourier expansion methods

Fang & Oosterlee (2008) proposed a new numerical method for pricing European options, which is called the COS method. This method is based on the Fourier transform and the Fourier-cosine expansion, and is shown to have the exponential convergence rate and linear computational complexity. And now, this novel method has been considered by many authors to price various options [see e.g. Ding *et al* (2011b), Fang & Oosterlee (2009), and the references therein].

We recall that the price S_t of underlying asset follows an exponential Lévy model. Let $X_t = \log(S_t/K)$, where K is the strike price. Consider the price of European call in the form:

$$V_0(K) = e^{-rT} \mathbb{E}[(S_T - K)^+] = e^{-rT} K \int_0^\infty (e^x - 1) f_T(x) dx, \quad (16)$$

where $f_T(x)$ is the probability density of X_T under the risk-neutral probability. Let $\varphi_T(u)$ be the characteristic function of $f_T(x)$. The main ideas of the COS method are to choose two numbers a and b such that the truncated integral approximates the infinite integral very well, i.e.,

$$\tilde{\varphi}_T(u) = \int_a^b e^{iux} f_T(x) dx \approx \int_{-\infty}^\infty e^{iux} f_T(x) dx = \varphi_T(u), \quad (17)$$

and to consider the Fourier-cosine expansion of $f_T(x)$ in $[a, b]$:

$$f_T(x) = \frac{1}{2} A_0 + \sum_{n=1}^\infty A_n \cos\left(n\pi \frac{x-a}{b-a}\right), \quad x \in [a, b], \quad (18)$$

where

$$A_n = \frac{2}{b-a} \int_a^b \tilde{p}_T(x) \cos\left(n\pi \frac{x-a}{b-a}\right) dx, \quad n = 0, 1, 2, \dots$$

From (17) we have

$$A_n = \frac{2}{b-a} \operatorname{Re} \left\{ \tilde{\varphi}_T\left(\frac{n\pi}{b-a}\right) \exp\left(-i \frac{na\pi}{b-a}\right) \right\}, \quad n = 0, 1, 2, \dots$$

Then, we get an approximation of $f_T(x)$ in (18) by

$$f_T(x) \approx \frac{1}{2} \tilde{A}_0 + \sum_{n=1}^{N-1} \tilde{A}_n \cos\left(n\pi \frac{x-a}{b-a}\right), \quad x \in [a, b], \quad (19)$$

where

$$\tilde{A}_n = \frac{2}{b-a} \operatorname{Re} \left\{ \varphi_T\left(\frac{n\pi}{b-a}\right) \exp\left(-i \frac{na\pi}{b-a}\right) \right\} \approx A_n, \quad n = 0, 1, \dots, N-1$$

Now, substituting (19) into (16), we obtain an approximation of the option price:

$$V_0(K) \approx Ke^{-rT} \left\{ \frac{1}{2} \tilde{A}_0 (\Phi_0(0, b) - \Psi_0(0, b)) + \sum_{n=1}^{N-1} \tilde{A}_n (\Phi_n(0, b) - \Psi_n(0, b)) \right\}, \quad (20)$$

where $\Phi_n(0, b)$ and $\Psi_n(0, b)$ are two integrals given by:

$$\Phi_n(c, d) = \int_c^d e^x \cos\left(n\pi \frac{x-a}{b-a}\right) dx \quad \text{and} \quad \Psi_n(c, d) = \int_c^d \cos\left(n\pi \frac{x-a}{b-a}\right) dx,$$

for any $[c, d] \subset [a, b]$, which are analytically given by

$$\begin{aligned} \Phi_n(c, d) &= \frac{1}{1 + \left(\frac{n\pi}{b-a}\right)^2} \left[\cos\left(n\pi \frac{d-a}{b-a}\right) e^d - \cos\left(n\pi \frac{c-a}{b-a}\right) e^c \right. \\ &\quad \left. + \frac{n\pi}{b-a} \sin\left(n\pi \frac{d-a}{b-a}\right) e^d - \frac{n\pi}{b-a} \sin\left(n\pi \frac{c-a}{b-a}\right) e^c \right], \\ \Psi_n(c, d) &= \begin{cases} \left[\sin\left(n\pi \frac{d-a}{b-a}\right) - \sin\left(n\pi \frac{c-a}{b-a}\right) \right] \frac{b-a}{n\pi} & n \neq 0, \\ (d-c) & n = 0 \end{cases}. \end{aligned}$$

Fang & Oosterlee (2008) also showed that, in most cases, the convergence rate of the COS formula (20) is exponential and the computational complexity is linear. They also discussed the truncation range for COS method, and gave a general formula to determine the interval of integration $[a, b]$ in that paper, which is given by

$$[a, b] = \left[c_1 - \delta \sqrt{c_2 + \sqrt{c_4}}, c_1 + \delta \sqrt{c_2 + \sqrt{c_4}} \right], \tag{21}$$

where c_1, c_2 , and c_4 are the first, second, and fourth cumulates of $X_t = \log(S_t/K)$. Also, the constant δ depends on the tolerance level in the approximation (17), and usually we choose $\delta = 10$. Meanwhile, Ding & U (2011) respectively applied the Fourier-sine and Fourier expansions to substitute the Fourier-cosine expansion in (18). A comprehensive analysis with numerical comparisons for these different methods is also given in their paper.

4. Fourier transform methods for pricing path dependent options

4.1 The CONV method for pricing Bermudan barrier options

Pricing Bermudan or barrier options is much harder than pricing European options. Because these options are depended on paths of the price process for the underlying assets. Recently, some new numerical integration methods based on Fourier transforms are proposed. Lord *et al* (2008) proposed an efficient and accurate FFT-based method, called the CONV method, to price Bermudan options under exponential Lévy models.

In the following we apply the CONV method to price a Bermudan barrier option in which the monitored dates may be many times more than the exercise dates. Denote

$$G(S) = \begin{cases} (S - K)^+, & \text{for call option,} \\ (K - S)^+, & \text{for put option,} \end{cases}$$

where K is the strike price, S is the spot price of the underlying asset. Let

$$\begin{cases} \mathbb{T} = \{t_{mL+l} : m = 0, 1, \dots, M, l = 0, 1, \dots, L-1\}, \\ \mathbb{T}_e = \{t_{mL} : m = 1, \dots, M-1\} \subset \mathbb{T}, \end{cases} \tag{22}$$

be the set of pre-specified monitored dates and the set of pre-specified exercise dates, respectively, before maturity T , where

$$0 = t_0 < t_1 < \dots < t_{ML} = T \quad \text{with} \quad \Delta t = t_k - t_{k-1} = T/(ML).$$

Consider a discrete American barrier option, which is monitored at every $t_k \in \mathbb{T}$ and can be exercised at each $t_k \in \mathbb{T}_e$, namely Bermudan barrier option, whose payoff is given by

$$G(S_{t_k})1_{\{S_{t_k} < H\}} + R_0 1_{\{S_{t_k} \geq H\}}, \quad t_k \in \mathbb{T}.$$

Here S_{t_k} is the price of the underlying asset at time $t_k \in \mathbb{T}$, $H > K$ is the constant barrier and R_0 is the contractual rebate. That is, this Bermudan barrier option is an up-and-out barrier option that cease to exist if the asset price S_{t_k} hits the barrier level H at one time $t_k \in \mathbb{T}$, and it can also be exercised at any time $t_k \in \mathbb{T}_e$ before maturity T .

Denote $V(S, t_k)$ the value of this Bermudan barrier option at time t_k and the spot price $S_{t_k} = S$. With help of the risk-neutral valuation formula, this price process can be computed recursively by the following backward induction:

$$\begin{cases} V(S, t_{ML}) = G(S)1_{\{S < H\}} + R_0 1_{\{S \geq H\}}, \\ C(S, t_k) = \mathbb{E}[e^{-r\Delta t} V(S_{t_{k+1}}, t_{k+1}) \mid S_{t_k} = S], \quad t_k \in \mathbb{T}, \\ V(S, t_k) = C(S, t_k)1_{\{S < H\}} + e^{-r(T-t_k)} R_0 1_{\{S \geq H\}}, \quad t_k \in \mathbb{T} \setminus \mathbb{T}_e, \\ V(S, t_k) = \max \{G(S), C(S, t_k)\} 1_{\{S < H\}} + e^{-r(T-t_k)} R_0 1_{\{S \geq H\}}, \quad t_k \in \mathbb{T}_e, \end{cases} \quad (23)$$

in specialty, the initial price is given

$$V(S, t_0) = C(S, t_0) = \mathbb{E}[e^{-r\Delta t} V(S, t_1) \mid S_{t_0} = S].$$

Here $r > 0$ is the interest rate, and $\mathbb{E}[\cdot \mid \cdot]$ is the conditional expectation under the risk-neutral probability.

Assume that the price process of the underlying asset is given by

$$S_t = S_0 e^{X_t}, \quad t \geq 0,$$

where X_t is a Lévy process and S_0 is the initial price. Let $f(\cdot \mid x)$ be the condition density of $X_{t_{k+1}}$ given $X_{t_k} = x$ for $t_k \in \mathbb{T}$. Set

$$g(x) = \begin{cases} (S_0 e^x - K)^+, & \text{for a call option,} \\ (K - S_0 e^x)^+, & \text{for a put option.} \end{cases}$$

Then the backward induction (23) can be rewritten by

$$\begin{cases} v(x, t_{ML}) = g(x)1_{\{x < h\}} + R_0 1_{\{x \geq h\}}, \\ c(x, t_k) = e^{-r\Delta t} \int_{-\infty}^{\infty} v(y, t_{k+1}) f(y \mid x) dy, \quad t_k \in \mathbb{T}, \\ v(x, t_k) = c(x, t_k)1_{\{x < h\}} + R_0 1_{\{x \geq h\}}, \quad t_k \in \mathbb{T} \setminus \mathbb{T}_e, \\ v(x, t_k) = \max \{g(x), c(x, t_k)\} 1_{\{x < h\}} + e^{-r(T-t_k)} R_0 1_{\{x \geq h\}}, \quad t_k \in \mathbb{T}_e, \end{cases} \quad (24)$$

and

$$v(x, t_0) = c(x, t_0) = e^{-r\Delta t} \int_{-\infty}^{\infty} v(y, t_1) f(y \mid x) dy,$$

where $v(x, t_k) = V(S_0 e^x, t_k)$ for any $t_k \in \mathbb{T}$, and $h = \log(H/S_0)$.

Since each Lévy process is stationary and has independent increments, the condition density $f(\cdot | x)$ possesses the property:

$$f(y|x) = f(y-x), \quad x, y \in \mathbb{R},$$

where $f(y)$ is the density of X_{t_1} under the initial condition $X_{t_0} = x$. Applying this property to infinite integrals $c(x, t_k)$ in (24) it becomes to

$$c(x, t_k) = e^{-r\Delta t} \int_{-\infty}^{\infty} v(y, t_{k+1}) f(y-x) dy = e^{-r\Delta t} \int_{-\infty}^{\infty} v(x+z, t_{k+1}) f(z) dz$$

for any $t_k \in \mathbb{T}$. Then, this integral can be rewritten as a convolution of $v(x+z, t_{k+1})$ and the function $f(-x)$, i.e.

$$c(x, t_k) = e^{-r\Delta t} f(-x) * v(x, t_{k+1}).$$

Thus we have

$$e^{\alpha x} c(x, t_k) = e^{-r\Delta t} [e^{\alpha x} f(-x)] * [e^{\alpha x} v(x, t_{k+1})], \tag{25}$$

where $\alpha > 0$ is chosen to improve the integrability. Now, the main idea of CONV method is that, taking the Fourier transform in the both sides of (25) and applying the Property P3 (Convolution), the integral becomes

$$\begin{aligned} e^{r\Delta t} \mathcal{F}[e^{\alpha x} c(x, t_k)](u) &= \mathcal{F}\{[e^{\alpha x} f(-x)] * [e^{\alpha x} v(x, t_{k+1})]\}(u) \\ &= \mathcal{F}[e^{\alpha x} f(-x)](u) \cdot \mathcal{F}[e^{\alpha x} v(x, t_{k+1})](u). \end{aligned}$$

Denote $\varphi(z)$ is the characteristic function of the density $f(x)$ on the complex plan \mathbb{C} . Then, by a simple calculation, we have

$$e^{r\Delta t} \mathcal{F}[e^{\alpha x} c(x, t_k)](u) = \varphi(- (u - i\alpha)) \cdot \mathcal{F}[e^{\alpha x} v(x, t_{k+1})](u).$$

Thus, taking the inverse Fourier transform we obtain

$$e^{\alpha x} c(x, t_k) = e^{-r\Delta t} \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-iux} \varphi(- (u - i\alpha)) \int_{-\infty}^{\infty} e^{iuy + \alpha y} v(y, t_{k+1}) dy du. \tag{26}$$

Denote

$$\kappa = \delta \sqrt{-\frac{\partial^2 \varphi_T(u)}{\partial u^2} \Big|_{u=0} + \left(\frac{\partial \varphi_T(u)}{\partial u} \Big|_{u=0}\right)^2},$$

where $\varphi_T(u)$ is the characteristic function of X_T , and δ is a proportionality constant. According to the suggestion from Lord *et al* (2008), we can take $\delta = 20$ for the GBM model and $\delta = 40$ for other exponential Lévy models. Let N be a power of 2. We consider the grid points on x -axes:

$$x_j = (j - \frac{1}{2}N)\Delta x, \quad j = 0, 1, \dots, N-1,$$

where $\Delta x = \kappa/N$. Furthermore, we also consider the grid points for the numerical integrals in (26):

$$u_j = (j - \frac{1}{2}N)\Delta u, \quad y_j = x_j, \quad j = 0, 1, \dots, N-1,$$

where $\Delta u = 2\pi/\kappa$. It is clear that these grids satisfy the Nyquist relation: $\Delta u \Delta y = 2\pi/N$. Now, for each $t_k \in \mathbb{T}$ and each $p = 0, 1, \dots, N-1$, approximating the integral in (26) with composite trapezoidal rule and the second integral with left rectangle rule yields

$$c(x_p, t_k) \approx e^{-\alpha x_p - r \Delta t} \frac{\Delta u \Delta y}{2\pi} \sum_{j=0}^{N-1} \left(e^{-iu_j x_p} \varphi(-(u_j - i\alpha)) \sum_{n=0}^{N-1} \omega_n e^{iu_j y_n + \alpha y_n} v(y_n, t_{k+1}) \right)$$

where the weights ω_n are chosen as $\omega_0 = \omega_{N-1} = \frac{1}{2}$ and $\omega_n = 1$ for $n = 1, \dots, N-2$. Noting that $u_0 = -\frac{1}{2}N\Delta u$ and $\Delta x = \Delta y = 2\pi/(N\Delta u)$, we have

$$c(x_p, t_k) \approx e^{-\alpha x_p - r \Delta t} e^{iu_0(y_0 - x_0)} (-1)^p \sum_{j=0}^{N-1} \left(e^{-ijp \frac{2\pi}{N}} e^{ij(y_0 - x_0)\Delta u} \varphi(-(u_j - i\alpha)) \right. \\ \left. \cdot \frac{1}{N} \sum_{n=0}^{N-1} e^{ijn \frac{2\pi}{N}} (-1)^n \omega_n e^{\alpha y_n} v(y_n, t_{k+1}) \right). \tag{27}$$

Now, we can employ the FFT algorithm to calculate the summations in the right side of (27). Once the integral $c(x_p, t_k)$ is computed, we can determine the early-exercise price $S_{t_k}^*$, $t_k \in \mathbb{T}_e$, by the procedure: for each $t_k \in \mathbb{T}_e$, locate j_k such that

$$c(x_{j_k}, t_k) - g(x_{j_k}) = 0, \tag{28}$$

or,

$$(c(x_{j_k}, t_k) - g(x_{j_k})) (c(x_{j_k+1}, t_k) - g(x_{j_k+1})) \leq 0. \tag{29}$$

In the case (28) set $x^*(t_k) = x_{j_k}$, and in the case (29) set $x^*(t_k) = \frac{1}{2}(x_{j_k} + x_{j_k+1})$. Then the early-exercise price at every $t_k \in \mathbb{T}_e$ is given by $S_{t_k}^* = S_0 e^{x^*(t_k)}$. Ding *et al* (2011b) gave a detail algorithm, which summarizes the above procedure, for pricing an up-and-out Bermudan barrier option, and the corresponding numerical experiments.

4.2 The COS method for pricing Bermudan barrier options

Recently, Fang & Oosterlee (2009) extended their COS method to price discrete early-exercise options under exponential Lévy models, and Fang & Oosterlee (2011) further considered such pricing problems under Heston’s model.

Assume that the price process of the underlying asset S_t follows an exponential Lévy model, and \mathbb{T} and \mathbb{T}_e are the set of pre-specified monitored dates and the set of pre-specified exercise dates, respectively, before the maturity T , which are defined by (22). In the following, we apply the COS method to price the Bermudan barrier option which defined in preceding subsection, whose payoff is given by

$$G(S_{t_k})1_{\{S_{t_k} < H\}} + R_0 1_{\{S_{t_k} \geq H\}}, \quad t_k \in \mathbb{T},$$

where $H > K$ is the constant barrier and R_0 is the contractual rebate. Denote $V(S, t_k)$, $t_k \in \mathbb{T}_e$, the value of this Bermudan barrier option at time t_k and the spot price $S_{t_k} = S$. As in the preceding subsection, with help of the risk-neutral valuation formula, this price process can be computed recursively by the backward induction (23). In specialty, the initial price is given by

$$V(S, t_0) = C(S, t_0) = \mathbb{E}[e^{-r\Delta t} V(S, t_1) \mid S_{t_0} = S]. \tag{30}$$

Here S is the spot price of underlying asset, $r > 0$ is the interest rate. Let $X_t = \log(S_t/K)$ be the logarithm of the underlying asset price S_t over the strike price K , and denote $x = \log(S/K)$ and $h = \log(H/K)$. Let $f(\cdot | x)$ be the condition density of $X_{t_{k+1}}$ given $X_{t_k} = x$ for $t_k \in \mathbb{T}$. Set

$$g(x) = \begin{cases} K(e^x - 1)^+, & \text{for a call option,} \\ K(1 - e^x)^+, & \text{for a put option.} \end{cases}$$

Then the backward induction (23) and the price formula (30) can be rewritten by

$$\begin{cases} v(x, t_{ML}) = g(x)1_{\{x < h\}} + R_0 1_{\{x \geq h\}}, \\ c(x, t_k) = e^{-r\Delta t} \int_{-\infty}^{\infty} v(y, t_{k+1})f(y | x)dy, & t_k \in \mathbb{T}, \\ v(x, t_k) = c(x, t_k)1_{\{x < h\}} + R_0 1_{\{x \geq h\}}, & t_k \in \mathbb{T} \setminus \mathbb{T}_e, \\ v(x, t_k) = \max \{g(x), c(x, t_k)\}1_{\{x < h\}} + e^{-r(T-t_k)}R_0 1_{\{x \geq h\}}, & t_k \in \mathbb{T}_e, \end{cases} \tag{31}$$

and

$$v(x, t_0) = c(x, t_0) = e^{-r\Delta t} \int_{-\infty}^{\infty} v(y, t_1)f(y | x)dy, \tag{32}$$

where $v(x, t_k) = V(Ke^x, t_k)$ for any $t_k \in \mathbb{T}$.

We consider the infinite integrals $c(x, t_k)$ in (31). Since $f(y|x)$ decays to zero very quickly as $y \rightarrow \pm\infty$ we may choose two bounds a and b , which can be selected by (21), such that

$$\bar{c}(x, t_k) = e^{-r\Delta t} \int_a^b v(y, t_{m+1})f(y | x)dy \approx c(x, t_k), \quad t_k \in \mathbb{T}. \tag{33}$$

without losing some significant accuracy. Note that the density $f(y | x)$ has the following Fourier-cosine expansion on $[a, b]$:

$$f(y | x) = \frac{2}{b-a} \sum_{j=0}^{\infty} \left[w_j \cos \left(j\pi \frac{y-a}{b-a} \right) \int_a^b f(u | x) \cos \left(j\pi \frac{u-a}{b-a} \right) du \right],$$

where $w_0 = \frac{1}{2}$ and $w_j = 1$ for all $j = 1, 2, 3, \dots$. We substitute this expansion into the integral (33) and then we can rewrite it by

$$\bar{c}(x, t_k) = \frac{(b-a)}{2} e^{-r\Delta t} \sum_{j=0}^{\infty} w_j F_j(x) V_j(t_{k+1}), \tag{34}$$

where for each $j = 1, 2, 3, \dots$,

$$V_j(t_{k+1}) = \frac{2}{b-a} \int_a^b v(y, t_{k+1}) \cos \left(j\pi \frac{y-a}{b-a} \right) dy, \tag{35}$$

and

$$F_j(x) = \frac{2}{b-a} \int_a^b f(y | x) \cos \left(j\pi \frac{y-a}{b-a} \right) dy. \tag{36}$$

Since the Fourier-cosine expansion has a high accuracy with a few terms, we can truncate the infinity series (34) and approximate $\bar{c}(x, t_k)$ by leaving the first N terms, i.e.,

$$\hat{c}(x, t_k) = \frac{(b-a)}{2} e^{-r\Delta t} \sum_{j=0}^{N-1} w_j F_j(x) V_j(t_{k+1}) \approx \bar{c}(x, t_k), \tag{37}$$

for any $t_k \in \mathbb{T}$, where $w_0 = w_{N-1} = \frac{1}{2}$ and $w_j = 1$ for $j = 1, \dots, N-2$. On the other hand, we can represent each integral (36) by approximation as the following

$$F_j(x) = \frac{2}{b-a} \operatorname{Re} \left\{ \exp \left(-i \frac{ja\pi}{b-a} \right) \int_a^b f(u | x) \exp \left(i \frac{ju\pi}{b-a} \right) du \right\} \\ \approx \frac{2}{b-a} \operatorname{Re} \left\{ \exp \left(-i \frac{ja\pi}{b-a} \right) \int_{-\infty}^{\infty} f(u | x) \exp \left(i \frac{j\pi}{b-a} u \right) du \right\}.$$

Let $\varphi(u; x)$ be the characteristic function of $f(\cdot | x)$. Then, we can approximate each $F_j(x)$ by

$$F_j(x) \approx \frac{2}{b-a} \operatorname{Re} \left\{ \exp \left(-i \frac{ja\pi}{b-a} \right) \cdot \varphi \left(\frac{j\pi}{b-a}; x \right) \right\}.$$

And so, we get the further numerical approximations of the integrals $\hat{c}(x, t_k)$ in (37) by

$$\tilde{c}(x, t_k) = e^{-r\Delta t} \sum_{j=0}^{N-1} w_j \operatorname{Re} \left\{ \exp \left(-i \frac{ja\pi}{b-a} \right) \cdot \varphi \left(\frac{j\pi}{b-a}; x \right) \right\} V_j(t_{k+1}) \approx \hat{c}(x, t_j). \quad (38)$$

In special, the approximation of initial price $v(x, t_0)$ in (32) is given by

$$\tilde{v}(x, t_0) = e^{-r\Delta t} \sum_{j=0}^{N-1} w_j \operatorname{Re} \left\{ \exp \left(-i \frac{ja\pi}{b-a} \right) \cdot \varphi \left(\frac{j\pi}{b-a}; x \right) \right\} V_j(t_1). \quad (39)$$

Meanwhile, since the characteristic function $\varphi(u; x)$ possesses the property:

$$\varphi(u; x) = \varphi(u) \cdot e^{iux}, \quad u \in \mathbb{R},$$

where $\varphi(u) = \varphi(u; 0)$, the approximations (38) and (39) can be simplified to

$$\tilde{c}(x, t_k) = e^{-r\Delta t} \sum_{j=0}^{N-1} w_j \operatorname{Re} \left\{ \exp \left(ij\pi \frac{x-a}{b-a} \right) \cdot \varphi \left(\frac{j\pi}{b-a} \right) \right\} V_j(t_{k+1}), \quad (40)$$

and the initial price of option

$$\tilde{v}(x, t_0) = e^{-r\Delta t} \sum_{j=0}^{N-1} w_j \operatorname{Re} \left\{ \exp \left(ij\pi \frac{x-a}{b-a} \right) \cdot \varphi \left(\frac{j\pi}{b-a} \right) \right\} V_j(t_1). \quad (41)$$

In order to use this approximate formulation we still need to compute the integrals $V_j(t_k)$. For convenience we introduce the following notions: for any $a \leq x_1 \leq x_2 \leq b$,

$$C_j(x_1, x_2; t_k) = \frac{2}{b-a} \int_{x_1}^{x_2} \tilde{c}(x, t_k) \cos \left(j\pi \frac{x-a}{b-a} \right) dx, \quad j = 0, 1, \dots, N-1,$$

and

$$D_j(x_1, x_2) = \frac{2R_0}{b-a} \int_{x_1}^{x_2} \cos \left(j\pi \frac{x-a}{b-a} \right) dx, \quad j = 0, 1, \dots, N-1,$$

where $\tilde{c}(x, t_{ML}) = g(x)$, and $\tilde{c}(x, t_k), t_k \in \mathbb{T}$, are given in (40).

We first calculate $V_j(t_{ML}), j = 0, 1, \dots, N-1$. We have

$$V_j(t_{ML}) = \begin{cases} C_j(0, h; t_{ML}) + D_j(h, b), & \text{for a call option,} \\ C_j(a, 0; t_{ML}) + D_j(h, b), & \text{for a put option,} \end{cases}$$

for all $j = 0, 1, \dots, N-1$. Denote

$$\Phi_j(x_1, x_2) = \int_{x_1}^{x_2} e^x \cos\left(j\pi \frac{x-a}{b-a}\right) dx \quad \text{and} \quad \Psi_k(x_1, x_2) = \int_{x_1}^{x_2} \cos\left(j\pi \frac{x-a}{b-a}\right) dx,$$

for any $a \leq x_1 \leq x_2 \leq b$ and $j = 0, \dots, N-1$. Then, by simple integration, these integrals admit the following analytic solutions:

$$\begin{aligned} \Phi_j(x_1, x_2) = & \frac{1}{1 + \left(\frac{j\pi}{b-a}\right)^2} \left[\cos\left(j\pi \frac{x_2-a}{b-a}\right) e^{x_2} - \cos\left(j\pi \frac{x_1-a}{b-a}\right) e^{x_1} \right. \\ & \left. + \frac{j\pi}{b-a} \sin\left(j\pi \frac{x_2-a}{b-a}\right) e^{x_2} - \frac{j\pi}{b-a} \sin\left(j\pi \frac{x_1-a}{b-a}\right) e^{x_1} \right], \end{aligned}$$

and

$$\Psi_j(x_1, x_2) = \left[\sin\left(j\pi \frac{x_2-a}{b-a}\right) - \sin\left(j\pi \frac{x_1-a}{b-a}\right) \right] \frac{b-a}{j\pi},$$

for $j = 0, \dots, N-1$, with $\Psi_0(x_1, x_2) = x_2 - x_1$. Moreover, by a simple calculation, $j = 0, \dots, N-1$, we have the following result:

$$D_j(x_1, x_2) = \frac{2R_0}{b-a} \Psi_j(x_1, x_2), \tag{42}$$

$$C_j(x_1, x_2; t_{ML}) = \frac{2}{b-a} \alpha K (\Phi_j(x_1, x_2) - \Psi_j(x_1, x_2)), \tag{43}$$

where α is a parameter such that $\alpha = 1$ for a call option and $\alpha = -1$ for a put option.

And next, we consider to calculate the integrals $V_j(t_k)$ for $t_k \in \mathbb{T} \setminus \mathbb{T}_e$. We have

$$V_j(t_k) = C_j(a, h; t_k) + e^{-r(T-t_{k-1})} D_j(h, b), \quad j = 0, 1, \dots, N-1.$$

Since the integral $D_j(x_1, x_2)$ has the analytic representation (45), we only need to calculate the integral $C_j(x_1, x_2; t_k)$. Fang & Oosterlee (2009) gave an efficient numerical algorithm which approximates $C_j(x_1, x_2; t_k)$ by using FFT method with the operation cost $O(N \log_2(N))$.

Finally, we consider to calculate the integrals $V_k(t_k)$ for $t_k \in \mathbb{T}_e$. It is clear that we should find the value $\tilde{v}(x, t_k)$ in the last equation in (31), or equivalently, to determine the early-exercise point x_k^* at each time t_k , which is the point where the continuation value is equal to the payoff, i.e., $\tilde{c}(x_k^*, t_k) = g(x_k^*)$. Let

$$h_k(y) = \tilde{c}(y, t_k) - g(y), \quad t_k \in \mathbb{T}_e.$$

Then, the problem becomes to find the root x_k^* of each equation $h_k(y) = 0$. Note that the function $\tilde{c}(y, t_k)$, which is given in (40), is bounded and smooth, and the function $g(y)$ is smooth except for $y = 0$ and bounded in $[a, b]$. We can use the Newton's method or the secant method to find the root x_k^* . Here if x_k^* is not in the interval $[a, b]$, we set it in the nearest boundary point a or b . Once we find the early-exercise point $x_k^*, t_k \in \mathbb{T}_e$, we have two different cases for an up-and-out barrier option:

Case 1: $x_k^* < h$, which means the early-exercise point doesn't hit the up barrier. Thus, We have the authority to decide to execute the option now or reserve it to the next time point. So we can split the integral that defines $V_j(t_k)$ into three parts: $[a, x_k^*]$, (x_k^*, h) and $[h, b]$. We have

$$V_j(t_k) = \begin{cases} C_j(a, x_k^*; t_k) + C_j(x_k^*, h; t_{ML}) + e^{-r(T-t_{k-1})}D_j(h, b), & \text{for a call option,} \\ C_j(a, x_k^*; t_{ML}) + C_j(x_k^*, h; t_k) + e^{-r(T-t_{k-1})}D_j(h, b), & \text{for a put option.} \end{cases}$$

Case 2: $x_k^* \geq h$, which means the early-exercise point hits the up barrier. Thus, the option integral can be split into two parts: $[a, h)$ and $[h, b]$:

$$V_j(t_k) = \begin{cases} C_j(a, h; t_k) + e^{-r(T-t_{k-1})}D_j(h, b), & \text{for a call option,} \\ C_j(a, h; t_{ML}) + e^{-r(T-t_{k-1})}D_j(h, b), & \text{for a put option,} \end{cases}$$

Ding *et al* (2011a) gave a detail algorithm, which summarizes the above procedure, for pricing an up-and-out Bermudan barrier option, and the corresponding numerical experiments.

4.3 The fast Hilbert transform approach for pricing barrier options

Feng & Linetsky (2008) presented a new numerical method to price discretely monitored barrier options under exponential Lévy models. Their method involves the relation with Hilbert transform (Property P4) and the Sine expansion in Hardy spaces. They also gave an efficient computational algorithm based on the fast Hilbert transform.

Let $\mathbb{T} = \{t_k : k = 1, \dots, M\}$ be the set of pre-specified monitored dates, where

$$0 = t_0 < t_1 < \dots < t_M = T \quad \text{with} \quad \Delta t = t_k - t_{k-1} = T/M.$$

We consider a European-type barrier put option whose payoff at maturity T is given by

$$G = 1_{\{S_{t_1} > L\}} 1_{\{S_{t_2} > L\}} \dots 1_{\{S_{t_M} > L\}} (K - S_T)^+,$$

where K is the strike price and $0 < L < K$ is the lower barrier. We also assume that the underlying asset price process is given by $S_t = Ke^{X_t}$ with $S_0 = S$, where X_t is a Lévy process started at $x = \log(S/K)$. Denoting $l = \log(L/K)$, then, with help of the risk-neutral valuation formula, the price of this option is given by

$$V(x, t_0) = e^{-rT} \mathbb{E} [1_{\{X_{t_1} > l\}} 1_{\{X_{t_2} > l\}} \dots 1_{\{X_{t_M} > l\}} K(1 - e^{X_{t_M}})^+ \mid X_0 = x],$$

which can be computed recursively by the following backward induction:

$$\begin{cases} V(x, t_M) = 1_{\{x > l\}} K(1 - e^x)^+, \\ V(x, t_k) = e^{-r\Delta t} 1_{\{x > l\}} \mathbb{E}[V(X_{t_{k+1}}, t_{k+1}) \mid X_{t_k} = x], \quad k = M-1, \dots, 1, \\ V(x, t_0) = e^{-rT} \mathbb{E}[e^{-r\Delta t} V(X_{t_1}, t_1) \mid X_{t_0} = x]. \end{cases} \tag{44}$$

Since each Lévy process is stationary and has independent increments, for each $k = 1, \dots, M-1$, we have

$$\mathbb{E}[V(X_{t_{k+1}}, t_{k+1}) \mid X_{t_k} = x] = \mathbb{E}[V(X_{t_1}, t_{k+1}) \mid X_{t_0} = x].$$

Thus, letting $P_{t_1}v(x) = \mathbb{E}[v(X_{t_1}) \mid X_{t_0} = x]$ be the expectation operator, from the backward induction (44) we get

$$\begin{cases} v_M(x) = 1_{\{x>l\}} \cdot K(1 - e^x)^+, \\ v_k(x) = 1_{\{x>l\}} P_{t_1}v_{k+1}(x), \quad j = M-1, \dots, 1, \\ v_0(x) = P_{t_1}v_1(x), \end{cases} \tag{45}$$

where $v_k(x) = e^{r\Delta t}V(x, t_k)$. And hence, the problem now becomes to find the function $v_0(x)$ from the backward induction (45).

Note that $1_{\{x>0\}} = \frac{1}{2}(1 + \text{sgn}(x))$ for all $x \in \mathbb{R}$. Using the Property P4 (Relation with Hilbert transform) we obtain

$$\mathcal{F}(1_{\{x>0\}} \cdot v)(u) = \frac{1}{2}(\mathcal{F}v(u) + i\mathcal{H}(\mathcal{F}v)(u)). \tag{46}$$

Let $a \in \mathbb{R}$ and \mathcal{T}_a be the transform operator: $\mathcal{T}_av(x) = v(x - a)$. Then, we have

$$1_{\{x>a\}} = \mathcal{T}_a1_{\{x>0\}} = \frac{1}{2}(1 + \mathcal{T}_a\text{sgn}(x)),$$

and hence,

$$1_{\{x>a\}} \cdot v(x) = \frac{1}{2}(v(x) + v(x) \cdot \mathcal{T}_a\text{sgn}(x)) = \frac{1}{2}(v(x) + \mathcal{T}_a(\text{sgn} \cdot \mathcal{T}_{-a}v)(x)).$$

Taking the Fourier transform to both sides of this equation, we have

$$\mathcal{F}(1_{\{x>a\}} \cdot v)(u) = \frac{1}{2}\mathcal{F}v(u) + \frac{1}{2}\mathcal{F}(\mathcal{T}_a(\text{sgn} \cdot \mathcal{T}_{-a}v))(u).$$

and using equation (46) we obtain

$$\mathcal{F}(1_{\{x>a\}} \cdot v)(u) = \frac{1}{2}\mathcal{F}v(u) + \frac{1}{2}ie^{iau}\mathcal{H}(e^{-ia y}\mathcal{F}v(y))(u). \tag{47}$$

On other hand, noting that, for each $k = 1, \dots, M-1$, the condition density of $X_{t_{k+1}}$ given $X_{t_k} = x$ possesses the property:

$$f(y|x) = f(y-x), \quad x, y \in \mathbb{R},$$

where $f(y)$ is the density of X_{t_1} under the initial condition $X_{t_0} = x$. The infinite integrals $P_{t_1}v_{k+1}(x)$ in (45) becomes to

$$P_{t_1}v_{k+1}(x) = \int_{-\infty}^{+\infty} v_{k+1}(y)f(y-x)dy = \int_{-\infty}^{+\infty} v_{k+1}(x+z)f(z)dz.$$

Then, this integral can be rewritten as a convolution of $v_{k+1}(x+z)$ and the function $f(-x)$, i.e.

$$P_{t_1}v_{k+1}(x) = f(-x) * v_{k+1}(x).$$

Note that $\text{supp}v_k(x) \subset (l, 0)$ for each $k = M, \dots, 1$ from the backward induction (45). We can take the Fourier transform in the both sides of (45). Applying the Property P3 (Convolution) we have

$$\mathcal{F}[P_{t_1}v_k(x)](u) = \mathcal{F}[f(-x) * v_{k+1}(x)](u) = \mathcal{F}[f(-x)](u) \cdot \mathcal{F}[v_{k+1}(x)](u).$$

Denote $\varphi(u)$ is the characteristic function of $f(x)$ on the complex plane \mathbb{C} . Then, by a simple calculation, we have

$$\mathcal{F}[P_{t_1} v_k(x)](u) = \varphi(-u) \cdot \mathcal{F}v_{k+1}(u). \quad (48)$$

Thus, using the formulas (47) and (48) we obtain

$$\begin{cases} \hat{v}_M(u) = \frac{K(1-e^{iu})}{iu} - \frac{K(1-e^{i(1+iu)})}{1+iu}, \\ \hat{v}_k(u) = \frac{1}{2}\varphi(-u)\hat{v}_{k+1}(u) + \frac{1}{2}ie^{iu} \mathcal{H}(e^{-iy} \varphi(-y)\hat{v}_{k+1}(y))(u), \\ \hat{v}_0(u) = \varphi(-u)\hat{v}_1(u). \end{cases} \quad k = M-1, \dots, 1, \quad (49)$$

Here $\hat{v}_k(u) = \mathcal{F}v_k(u)$ for each k . Applying the truncated Sinc approximation, Feng & Linetsky (2008) obtained the discretization of $\hat{v}_k(u)$:

$$\begin{aligned} \hat{v}_k(nh) &= \frac{1}{2}\varphi(-nh)\hat{v}_{k+1}(nh) \\ &+ \frac{ie^{inh}}{2\pi} \sum_{j=-N, j \neq n}^N e^{-ijhl} \varphi(-jh)\hat{v}_{k+1}(jh) \frac{1 - (-1)^{n-j}}{n-j}, \end{aligned} \quad (50)$$

for $n = -N, \dots, N$ and $k = M-1, \dots, 1$, where N is a positive integer and h is the discretization step size. Then, the function $v_0(x)$ can be computed by the discretised inversion Fourier transform:

$$v_0(x) = \frac{1}{2\pi} \sum_{j=-N}^N e^{-ijhx} \varphi(-jh)\hat{v}_1(jh)h.$$

Furthermore, Feng & Linetsky (2008) showed that the computation (50) involves a Toeplitz matrix-vector multiplication, which can be accomplished in $O(N \log_2 N)$ operations using the FFT technique. They also referred the corresponding algorithm of computing the discrete Hilbert transform via the FFT as the *Fast Hilbert Transform*.

5. References

- Barndorff-Nielsen, O. (1997). Normal inverse Gaussian distributions and stochastic volatility modelling, *Scandinavian Journal of Statistics*, Vol. 24, pp. 1–13.
- Bates, D. S. (1996). Jumps and stochastic volatility: exchange rate processes implicit in Deutsche mark options, *Rev. Financ. Stud.*, Vol. 9, pp. 69–107.
- Black, F. & Scholes, M. (1973). The pricing of options and corporate liabilities, *Journal of Political Economy*, Vol. 81 (No. 3), pp. 637–654.
- Borak, S., Detlefsen, K. & Härdle, W. (2005). FFT based option pricing, SFB 649 Discussion paper 2005-011. <http://sfb649.wiwi.hu-berlin.de>
- Carr, P., Geman, H., Madan, D. & Yor, M. (2002). The fine structure of asset returns: An empirical investigation. *Journal of Business*, Vol. 75, pp. 305–332.
- Carr, P. & Madan, D. (1999). Option valuation using the fast Fourier transform, *Journal of Computational Finance*, Vol. 2 pp. 61–73.
- Carr, P. & Wu, L. (2003). The finite moment log stable process and option pricing, *Journal of Finance*, Vol. 58, pp. 753–777.
- Cont, R. & Tankov, P. (2003). *Financial Modelling with Jump Processes*, Chapman & Hall/ CRC Press.

- Dempster, M. A. & Hong, S. S. G. (2002). Spread option valuation and the fast Fourier transform, *Mathematical Finance–Bachelier Congress 2000*, Spring, pp. 203–220.
- Ding, D & U, S. C. (2010). An accurate and stable FFT-based method for pricing options under exp-Levy processes, *The Proceedings of ISCM II - EPMSC XII*, American Institute of Physics, pp. 741–746.
- Ding, D & U, S. C. (2011). Efficient option pricing methods based on Fourier series expansions, *Journal of Mathematical Research and Exposition*, Vol. 31, pp. 12–22.
- Ding, D., Huang, H. & Zhao, J. (2011a). An efficient algorithm for Bermudan barrier option pricing, Working paper, University of Macau.
- Ding, D., Weng, Z. & Zhao, J. (2011b). Pricing Bermudan barrier options via a fast and accurate FFT-based Method, to appear in the Proceedings of CiSE 2011, IEEE.
- Duffie, D., Pan, J. & Singleton, K. (2000). Transform analysis and asset pricing for affine jump-diffusions, *Econometrica*, Vol. 68, pp. 1343–1376.
- Dufresne, D., Garrido, J. & Morales, M. (2009). Fourier inversion formulas in option pricing and insurance, *Methodol. Comput. Appl. Probab.*, Vol. 11, pp. 359–383.
- Eberlein, E., Glau, K., & Papapantoleon, A. (2010). Analysis of Fourier transform valuation formulas and application, *Applied Mathematical Finance*, Vol. 17, pp. 211–240.
- Fang, F. & Oosterlee, C. W. (2008). A novel pricing method for European option based on Fourier-cosine series expansions, *SIAM J. Sci. Comput.*, Vol. 31, pp. 826–848.
- Fang, F. & Oosterlee, C. W. (2009). Pricing early-exercise and discrete barrier options by Fourier-cosine series expansions, *Numer. Math.*, Vol. 114, pp. 27–62.
- Fang, F. & Oosterlee, C. W. (2011). A Fourier-based valuation method for Bermudan and barrier options under Heston’s model, *SIAM J. Financial Math.*, Vol. 2, pp. 439–463.
- Feng, L. & Linetsky, V. (2008). Pricing discretely monitored barrier options and defaultable bonds in Lévy process models: a fast Hilbert transform approach. *Mathematical Finance*, Vol. 18, pp. 337–384.
- Gurland, J. (1948). Inversion formula for the distribution of ratios, *Ann. Math. Statist.*, Vol. 19, pp. 228–237.
- Heston, S. (1993). A closed-form solution for options with stochastic volatility with applications to bond and currency options, *Rev. Financ. Stud.*, Vol. 6, pp. 327–343.
- Hurd, T. & Zhou, Z. (2010). A Fourier transform method for spread option pricing, *SIAM J. Finan. Math.*, Vol. 1, pp. 142–157.
- Lee, R. (2004). Option pricing by transform methods: Extensions, unification, and error control, *Journal of Computational Finance*, Vol. 7, pp. 51–86.
- Lewis, A. (2001). A simple option formula for general jump-diffusion and other exponential Lévy processes. <http://www.optioncity.net>
- Lord, R., Fang, F., Bervoets, F. & Oosterlee, C. W. (2008). A fast and accurate FFT-based method for pricing early-exercise options under Lévy processes. *SIAM J. Sci. Comput.*, Vol. 30, pp. 1678–1705.
- Kou, S. G. (2002). A jump-diffusion model for option pricing, *Management Science*, Vol. 48, pp. 1086–1101.
- Kwok, Y. K., Leung, K. S. & Wong, H. Y. (2010). Efficient options pricing using the fast Fourier transform. <http://ssrn.com/abstract=1534544>
- Madan, D., Carr, P. & Chang, E. (1998). The variance gamma process and option pricing, *European Financial Review*, Vol. 2, pp. 79–105.
- Merton, R. (1973). Theory of rational option pricing, *Bell Journal of Economics and Management Science*, Vol. 4, pp. 141–183.

- Merton, R. (1976). Option pricing when underlying stock returns are discontinuous. *J. Financ. Econ.*, Vol. 3, pp. 125–144.
- Schmelzle, M. (2010). Option Pricing Formulae using Fourier Transform: Theory and Application. <http://pfadintegral.com/articles/option-pricing-formulae-using-fourier-transform/>
- Shephard, N. G. (1991). From characteristic function to distribution function: a simple framework for the theory, *Econometric Theory*, Vol. 7, pp. 519–529.
- Stein, E. & Stein, F. (1991). Stock price distribution with stochastic volatility: An analytic approach, *Rev. Financ. Stud.*, Vol. 4, pp. 727–752.

INTECH

INTECH



Fourier Transform Applications

Edited by Dr Salih Salih

ISBN 978-953-51-0518-3

Hard cover, 300 pages

Publisher InTech

Published online 25, April, 2012

Published in print edition April, 2012

The book focuses on Fourier transform applications in electromagnetic field and microwave, medical applications, error control coding, methods for option pricing, and Helbert transform application. It is hoped that this book will provide the background, reference and incentive to encourage further research and results in these fields as well as provide tools for practical applications. It provides an applications-oriented analysis written primarily for electrical engineers, control engineers, signal processing engineers, medical researchers, and the academic researchers. In addition the graduate students will also find it useful as a reference for their research activities.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Deng Ding (2012). Fourier Transform Methods for Option Pricing, Fourier Transform Applications, Dr Salih Salih (Ed.), ISBN: 978-953-51-0518-3, InTech, Available from: <http://www.intechopen.com/books/fourier-transform-applications/fourier-transform-methods-for-option-pricing>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821