


Triphenyl[(4-phenylbenzoyl)methyl]phosphonium trifluoromethanesulfonate

Corrado Rizzoli, Kazem Karami and Mina Mohamadi Salah

Acta Cryst. (2010). **E66**, o2675–o2676

This open-access article is distributed under the terms of the Creative Commons Attribution Licence
<http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals Online is available from journals.iucr.org

Triphenyl[(4-phenylbenzoyl)methyl]- phosphonium trifluoromethanesulfonate

Corrado Rizzoli,^{a*} Kazem Karami^b and Mina Mohamadi
 Salah^b

^aDipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, Università degli Studi di Parma, Viale G. P. Usberti 17/A, I-43100 Parma, Italy, and ^bDepartment of Chemistry, Isfahan University of Technology, Isfahan 84156/83111, Iran

Correspondence e-mail: corrado.rizzoli@unipr.it


Received 20 September 2010; accepted 24 September 2010

Key indicators: single-crystal X-ray study; $T = 294\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.048; wR factor = 0.139; data-to-parameter ratio = 13.8.

In the cation of the title compound, $\text{C}_{32}\text{H}_{26}\text{OP}^+\cdot\text{CF}_3\text{O}_3\text{S}^-$, the dihedral angle between the benzene rings of the biphenyl group is $42.37(8)^\circ$. In the crystal, the cations and anions interact through intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming chains parallel to the b axis. These chains are further linked by $\text{C}-\text{H}\cdots\pi$ stacking interactions into layers parallel to the bc plane.

Related literature

For the synthesis and characterization of phosphorus ylide metal complexes, see: Kalyanasundari *et al.* (1995, 1999); Laavanya *et al.* (2001); Vicente *et al.* (1985); Karami (2007, 2008); Akkurt *et al.* (2008). For related structures, see: Karami & Büyükgüngör (2009); Shao *et al.* (1982). For the synthesis of the title compound, see: Burmeister *et al.* (1973).



Experimental

Crystal data



$M_r = 606.58$

Monoclinic, $P2_1/c$

$a = 9.0559(10)\text{ \AA}$

$b = 19.382(2)\text{ \AA}$

$c = 16.5396(19)\text{ \AA}$

$\beta = 92.577(2)^\circ$

$V = 2900.1(6)\text{ \AA}^3$

$Z = 4$

$\text{Mo K}\alpha$ radiation

$\mu = 0.22\text{ mm}^{-1}$

$T = 294\text{ K}$

$0.25 \times 0.20 \times 0.17\text{ mm}$

Data collection

Bruker SMART 1000 CCD

diffractometer

Absorption correction: multi-scan
 (SADABS; Bruker, 1998)

$T_{\min} = 0.936$, $T_{\max} = 0.974$

29195 measured reflections

5245 independent reflections

3748 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$

$wR(F^2) = 0.139$

$S = 1.06$

5245 reflections

379 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 0.56\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$, $Cg2$ and $Cg3$ are the centroids of the C1–C6, C21–C26 and C15–C20 phenyl rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1–H1B \cdots O4 ⁱ	0.97	2.22	3.191 (4)	178
C26–H26 \cdots O3 ⁱⁱ	0.93	2.45	3.373 (3)	174
C32–H32 \cdots O3 ⁱⁱ	0.93	2.46	3.369 (4)	168
C1–H1A \cdots Cg1 ⁱⁱⁱ	0.97	2.84	3.780 (3)	164
C10–H10 \cdots Cg2 ^{iv}	0.93	3.02	3.767 (4)	138
C23–H23 \cdots Cg3 ^v	0.93	2.91	3.788 (4)	159

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x, -y + 1, -z$; (iv) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (v) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT-Plus (Bruker, 1998); data reduction: SAINT-Plus; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHEXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and SCHAKAL97 (Keller, 1997); software used to prepare material for publication: SHEXL97 and PARST95 (Nardelli, 1995).

Financial support from the Università degli Studi di Parma is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2176).

References

- Akkurt, M., Karami, K., Yalçın, Ş. P. & Büyükgüngör, O. (2008). *Acta Cryst. E* **64**, m612–m613.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Molteri, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bruker (1998). SMART, SAINT-Plus and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burmeister, J. L., Silver, J. L., Weleski, E. T., Schweizer, E. E. & Kopay, C. M. (1973). *Synth. Inorg. Met. Org. Chem.* **3**, 339–358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Kalyanasundari, M., Panchantheswaran, K., Parthasarathi, V. & Robinson, W. T. (1999). *Bull. Chem. Soc. Jpn.* **72**, 33–36.
- Kalyanasundari, M., Panchantheswaran, K., Robinson, W. T. & Wen, H. (1995). *J. Organomet. Chem.* **491**, 103–109.
- Karami, K. (2007). *J. Chem. Res. (S)*, pp. 725–727.
- Karami, K. (2008). *Transition Met. Chem.* **33**, 819–823.
- Karami, K. & Büyükgüngör, O. (2009). *Acta Cryst. E* **65**, o296.
- Keller, E. (1997). SCHAKAL97. University of Freiburg, Germany.
- Laavanya, P., Venkatasubramanian, U., Panchantheswaran, K. & Bauer, J. A. K. (2001). *Chem. Commun.* pp. 1660–1661.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.

organic compounds

- Shao, M., Jin, X., Tang, Y. & Huang, Q. Y. (1982). *Tetrahedron Lett.* **23**, 5343–5346.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Vicente, J., Chicote, M. T., Cayuelas, J. A., Fernandez-Baeza, J., Jones, P. G., Sheldrick, G. M. & Espinet, P. (1985). *J. Chem. Soc. Dalton Trans.* pp. 1163–1168.

supplementary materials

Acta Cryst. (2010). E66, o2675–o2676 [doi:10.1107/S16005368100038286]

Triphenyl[(4-phenylbenzoyl)methyl]phosphonium trifluoromethanesulfonate

C. Rizzoli, K. Karami and M. M. Salah

Comment

The synthesis of α -ketostabilized phosphorus ylides has attracted much interest over the last decades due to their stability and ability to act as bifunctional ligands (C- versus O-coordination; Kalyanasundari *et al.*, 1995; Kalyanasundari *et al.*, 1999; Laavanya *et al.*, 2001; Vicente *et al.*, 1985). In this respect, the preferred coordination modes of 4-fluorobenzoyloxy-methylenetriphenylphosphorane ylide (FBPPY), 4-chlorobenzyl-oxymethylenetriphenylphosphorane ylide (CBPPY), and 4-methoxybenzoylmethylenetriphenylphosphorane ylide (MOBPPY) to transition metals such as mercury (II), silver (I) and palladium (II) have been recently investigated by our group (Karami, 2007, 2008; Akkurt *et al.*, 2008). As a part of our ongoing study on the synthesis and characterization of new trifluoromethanesulfonate phosphonium ylides, the title compound has been prepared according to the sequence shown in Figure 3 (Burmeister *et al.*, 1973), and its crystal structure is reported herein.

In the title compound (Fig. 1), bond lengths and angles within the phosphonium cation are not unusual and comparable to those observed for the related 4-methoxybenzoylmethyl derivative (Karami & Büyükgüngör, 2009). The P1–C1 bond length (1.798 (2) Å) is significantly longer than that reported in the free ylide (1.711 Å) of formula $\text{Ph}_3\text{PC}(\text{H})\text{COPh}$ (Shao *et al.*, 1982). The dihedral angle between the benzene rings of the biphenyl group is 42.37 (8) $^\circ$. Unlike the 4-methoxybenzoylmethyl derivative, the conformation of the cation is not stabilized by intramolecular hydrogen bonds. In the crystal packing (Fig. 2), cations and anions are linked by intermolecular C—H \cdots O hydrogen bonds into chains running parallel to the *b* axis (see Fig. 2 & Table 1). The chains further interact through C—H \cdots π stacking interactions to form layers parallel to the *bc* plane (see Fig. 2 & Table 1).

Experimental

The title compound was obtained by reaction of (4-phenylbenzoylmethyl)triphenylphosphonium bromide and AgOTf (OTf : trifluoromethanesulfonate) in dry acetone in a 1:1 molar ratio under stirring for 12 h (Fig. 3). The precipitate obtained was washed several times with dry diethyl ether and dried in a vacuum. Orange crystals of the title compound suitable for X-ray analysis formed by addition of dry diethyl ether to a chloroform solution. The crystals are air stable and resistant against moisture.

Refinement

All H atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93–0.97 Å, and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 40% probability level. H atoms are presented as a small spheres of arbitrary radius.



Fig. 2. A view of the C—H···O and C—H···π interactions (dashed lines) in the crystal structure of the title compound.




Fig. 3. The procedure adopted for the synthesis of the title compound.

Triphenyl[(4-phenylbenzoyl)methyl]phosphonium trifluoromethanesulfonate

Crystal data

$C_{32}H_{26}OP^+\cdot CF_3O_3S^-$	$F(000) = 1256$
$M_r = 606.58$	$D_x = 1.389 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 744 reflections
$a = 9.0559 (10) \text{ \AA}$	$\theta = 6.3\text{--}22.4^\circ$
$b = 19.382 (2) \text{ \AA}$	$\mu = 0.22 \text{ mm}^{-1}$
$c = 16.5396 (19) \text{ \AA}$	$T = 294 \text{ K}$
$\beta = 92.577 (2)^\circ$	Block, orange
$V = 2900.1 (6) \text{ \AA}^3$	$0.25 \times 0.20 \times 0.17 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART 1000 CCD diffractometer	5245 independent reflections
Radiation source: fine-focus sealed tube graphite	3748 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.043$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 1998)	$\theta_{\text{max}} = 25.3^\circ, \theta_{\text{min}} = 1.6^\circ$
$T_{\text{min}} = 0.936, T_{\text{max}} = 0.974$	$h = -10 \rightarrow 10$
29195 measured reflections	$k = -23 \rightarrow 23$
	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: difference Fourier map
$wR(F^2) = 0.139$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.082P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
5245 reflections	$(\Delta/\sigma)_{\max} < 0.001$
379 parameters	$\Delta\rho_{\max} = 0.56 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$

Special details

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.05841 (6)	0.19921 (3)	0.17248 (4)	0.03733 (18)
S1	0.78232 (9)	0.42703 (4)	0.27665 (5)	0.0666 (2)
F1	0.5564 (3)	0.35241 (16)	0.3102 (3)	0.1880 (17)
F2	0.5301 (3)	0.45799 (15)	0.3327 (2)	0.1577 (13)
F3	0.6656 (4)	0.39779 (16)	0.41126 (19)	0.1544 (13)
O1	0.29798 (19)	0.28836 (9)	0.13868 (13)	0.0625 (5)
O2	0.7193 (4)	0.4436 (2)	0.19884 (16)	0.1420 (13)
O3	0.8484 (3)	0.48364 (11)	0.31695 (15)	0.0894 (7)
O4	0.8683 (3)	0.36544 (12)	0.28273 (18)	0.1039 (9)
C1	0.0380 (3)	0.28830 (11)	0.14394 (15)	0.0435 (6)
H1A	-0.0207	0.2913	0.0935	0.052*
H1B	-0.0145	0.3126	0.1851	0.052*
C2	0.1864 (3)	0.32291 (12)	0.13352 (15)	0.0449 (6)
C3	0.1899 (2)	0.39764 (12)	0.11621 (14)	0.0433 (6)
C4	0.0651 (3)	0.43896 (12)	0.11089 (15)	0.0459 (6)
H4	-0.0267	0.4202	0.1211	0.055*
C5	0.0757 (3)	0.50804 (12)	0.09048 (15)	0.0453 (6)
H5	-0.0088	0.5354	0.0883	0.054*
C6	0.2109 (3)	0.53701 (12)	0.07326 (14)	0.0433 (6)
C7	0.3360 (3)	0.49527 (13)	0.08168 (17)	0.0535 (7)
H7	0.4282	0.5140	0.0724	0.064*
C8	0.3263 (3)	0.42769 (13)	0.10323 (17)	0.0523 (7)
H8	0.4119	0.4013	0.1093	0.063*

supplementary materials

C9	0.2251 (3)	0.60970 (12)	0.04788 (14)	0.0437 (6)
C10	0.1499 (3)	0.66280 (13)	0.08446 (16)	0.0549 (7)
H10	0.0825	0.6526	0.1235	0.066*
C11	0.1740 (3)	0.73053 (14)	0.06349 (18)	0.0651 (8)
H11	0.1235	0.7657	0.0888	0.078*
C12	0.2723 (3)	0.74644 (14)	0.00535 (18)	0.0613 (7)
H12	0.2913	0.7923	-0.0070	0.074*
C13	0.3418 (3)	0.69483 (15)	-0.03417 (18)	0.0601 (7)
H13	0.4048	0.7055	-0.0753	0.072*
C14	0.3188 (3)	0.62676 (13)	-0.01322 (17)	0.0539 (7)
H14	0.3667	0.5919	-0.0404	0.065*
C15	0.1376 (3)	0.14886 (12)	0.09454 (14)	0.0414 (5)
C16	0.0457 (3)	0.12156 (13)	0.03274 (15)	0.0530 (6)
H16	-0.0555	0.1299	0.0320	0.064*
C17	0.1047 (4)	0.08214 (16)	-0.02744 (18)	0.0690 (8)
H17	0.0437	0.0642	-0.0690	0.083*
C18	0.2538 (4)	0.06968 (16)	-0.02533 (19)	0.0763 (9)
H18	0.2933	0.0427	-0.0655	0.092*
C19	0.3454 (3)	0.09620 (16)	0.03476 (19)	0.0677 (8)
H19	0.4463	0.0872	0.0353	0.081*
C20	0.2880 (3)	0.13635 (13)	0.09469 (15)	0.0511 (6)
H20	0.3505	0.1550	0.1352	0.061*
C21	0.1644 (2)	0.19295 (12)	0.26621 (14)	0.0398 (5)
C22	0.1840 (3)	0.24969 (14)	0.31672 (16)	0.0594 (7)
H22	0.1494	0.2929	0.3003	0.071*
C23	0.2550 (4)	0.24155 (19)	0.39123 (19)	0.0771 (9)
H23	0.2690	0.2796	0.4249	0.093*
C24	0.3053 (3)	0.1785 (2)	0.41654 (18)	0.0728 (9)
H24	0.3536	0.1739	0.4670	0.087*
C25	0.2845 (3)	0.12184 (17)	0.36745 (17)	0.0627 (7)
H25	0.3177	0.0788	0.3851	0.075*
C26	0.2145 (3)	0.12866 (13)	0.29199 (15)	0.0476 (6)
H26	0.2011	0.0903	0.2586	0.057*
C27	-0.1209 (2)	0.16470 (12)	0.19059 (13)	0.0389 (5)
C28	-0.2382 (3)	0.20647 (13)	0.20946 (15)	0.0473 (6)
H28	-0.2285	0.2542	0.2078	0.057*
C29	-0.3701 (3)	0.17695 (15)	0.23093 (18)	0.0591 (7)
H29	-0.4494	0.2050	0.2431	0.071*
C30	-0.3841 (3)	0.10673 (15)	0.23429 (17)	0.0576 (7)
H30	-0.4729	0.0872	0.2488	0.069*
C31	-0.2677 (3)	0.06509 (14)	0.21634 (16)	0.0555 (7)
H31	-0.2781	0.0174	0.2188	0.067*
C32	-0.1356 (3)	0.09315 (12)	0.19477 (15)	0.0469 (6)
H32	-0.0567	0.0647	0.1831	0.056*
C33	0.6237 (4)	0.4073 (2)	0.3325 (3)	0.0963 (13)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0344 (3)	0.0359 (3)	0.0418 (3)	-0.0018 (2)	0.0030 (2)	-0.0002 (3)
S1	0.0717 (5)	0.0581 (5)	0.0714 (5)	-0.0069 (4)	0.0203 (4)	-0.0080 (4)
F1	0.116 (2)	0.126 (2)	0.327 (5)	-0.0639 (19)	0.070 (3)	-0.014 (3)
F2	0.0997 (18)	0.150 (2)	0.229 (3)	0.0702 (17)	0.0710 (19)	0.072 (2)
F3	0.179 (3)	0.161 (3)	0.130 (2)	0.054 (2)	0.077 (2)	0.072 (2)
O1	0.0444 (10)	0.0477 (11)	0.0966 (15)	0.0018 (8)	0.0148 (10)	0.0106 (10)
O2	0.148 (3)	0.214 (4)	0.0626 (17)	-0.034 (2)	-0.0075 (17)	0.0103 (19)
O3	0.1034 (17)	0.0534 (12)	0.1110 (19)	-0.0142 (12)	0.0002 (14)	-0.0081 (12)
O4	0.1033 (18)	0.0602 (14)	0.153 (3)	0.0123 (13)	0.0578 (17)	-0.0148 (14)
C1	0.0414 (13)	0.0375 (13)	0.0520 (14)	-0.0040 (10)	0.0054 (11)	0.0025 (10)
C2	0.0416 (14)	0.0432 (13)	0.0507 (14)	-0.0023 (11)	0.0098 (11)	0.0003 (11)
C3	0.0420 (13)	0.0422 (13)	0.0462 (14)	-0.0041 (10)	0.0084 (10)	-0.0010 (10)
C4	0.0384 (13)	0.0478 (14)	0.0517 (14)	-0.0064 (11)	0.0058 (11)	0.0020 (11)
C5	0.0424 (13)	0.0436 (14)	0.0503 (14)	0.0012 (11)	0.0061 (11)	0.0007 (11)
C6	0.0464 (14)	0.0426 (13)	0.0414 (13)	-0.0025 (11)	0.0066 (10)	-0.0041 (10)
C7	0.0417 (14)	0.0457 (15)	0.0742 (18)	-0.0067 (11)	0.0147 (13)	0.0010 (13)
C8	0.0387 (13)	0.0423 (14)	0.0769 (19)	0.0013 (11)	0.0137 (12)	-0.0007 (12)
C9	0.0452 (13)	0.0424 (13)	0.0435 (13)	-0.0021 (11)	0.0018 (11)	-0.0002 (10)
C10	0.0666 (17)	0.0495 (16)	0.0497 (15)	0.0016 (13)	0.0129 (13)	-0.0033 (12)
C11	0.091 (2)	0.0433 (15)	0.0612 (18)	0.0113 (15)	0.0022 (16)	-0.0048 (13)
C12	0.0691 (18)	0.0428 (15)	0.0709 (19)	-0.0043 (13)	-0.0090 (15)	0.0096 (13)
C13	0.0476 (15)	0.0601 (18)	0.0730 (19)	-0.0016 (13)	0.0078 (13)	0.0194 (15)
C14	0.0504 (15)	0.0471 (15)	0.0652 (17)	0.0008 (12)	0.0148 (13)	0.0026 (12)
C15	0.0467 (14)	0.0397 (13)	0.0382 (13)	0.0008 (10)	0.0056 (10)	0.0023 (10)
C16	0.0560 (15)	0.0547 (16)	0.0481 (15)	-0.0007 (12)	0.0000 (12)	-0.0033 (12)
C17	0.088 (2)	0.068 (2)	0.0516 (17)	-0.0058 (17)	0.0021 (15)	-0.0153 (14)
C18	0.102 (3)	0.071 (2)	0.0576 (19)	0.0077 (19)	0.0256 (18)	-0.0164 (15)
C19	0.0650 (18)	0.0716 (19)	0.068 (2)	0.0138 (15)	0.0215 (15)	-0.0001 (16)
C20	0.0474 (15)	0.0577 (16)	0.0488 (15)	0.0050 (12)	0.0083 (12)	-0.0014 (12)
C21	0.0345 (12)	0.0438 (13)	0.0413 (13)	-0.0028 (10)	0.0044 (9)	-0.0023 (10)
C22	0.0720 (19)	0.0517 (16)	0.0539 (17)	-0.0014 (14)	-0.0032 (14)	-0.0094 (13)
C23	0.087 (2)	0.085 (2)	0.0588 (19)	-0.0074 (19)	-0.0087 (16)	-0.0219 (17)
C24	0.0621 (19)	0.107 (3)	0.0481 (17)	0.0019 (18)	-0.0105 (14)	-0.0032 (17)
C25	0.0531 (16)	0.078 (2)	0.0571 (18)	0.0112 (14)	-0.0008 (13)	0.0138 (15)
C26	0.0446 (14)	0.0495 (15)	0.0485 (14)	0.0015 (11)	0.0005 (11)	0.0010 (11)
C27	0.0351 (12)	0.0427 (13)	0.0387 (12)	-0.0022 (10)	-0.0001 (10)	0.0016 (10)
C28	0.0417 (13)	0.0434 (14)	0.0567 (15)	-0.0007 (11)	0.0018 (11)	-0.0025 (11)
C29	0.0372 (14)	0.0685 (19)	0.0719 (19)	0.0000 (13)	0.0076 (13)	-0.0009 (14)
C30	0.0399 (14)	0.0691 (19)	0.0642 (18)	-0.0146 (13)	0.0041 (12)	0.0028 (14)
C31	0.0545 (16)	0.0468 (15)	0.0649 (18)	-0.0150 (12)	0.0011 (13)	0.0038 (12)
C32	0.0434 (13)	0.0422 (14)	0.0554 (15)	-0.0022 (11)	0.0039 (11)	0.0027 (11)
C33	0.078 (2)	0.077 (3)	0.137 (4)	0.019 (2)	0.036 (2)	0.028 (2)

supplementary materials

Geometric parameters (\AA , $^\circ$)

P1—C21	1.790 (2)	C13—H13	0.9300
P1—C15	1.792 (2)	C14—H14	0.9300
P1—C27	1.794 (2)	C15—C20	1.383 (3)
P1—C1	1.798 (2)	C15—C16	1.393 (3)
S1—O3	1.404 (2)	C16—C17	1.381 (4)
S1—O2	1.421 (3)	C16—H16	0.9300
S1—O4	1.427 (2)	C17—C18	1.371 (4)
S1—C33	1.783 (4)	C17—H17	0.9300
F1—C33	1.273 (5)	C18—C19	1.366 (4)
F2—C33	1.298 (4)	C18—H18	0.9300
F3—C33	1.354 (5)	C19—C20	1.380 (4)
O1—C2	1.212 (3)	C19—H19	0.9300
C1—C2	1.519 (3)	C20—H20	0.9300
C1—H1A	0.9700	C21—C26	1.387 (3)
C1—H1B	0.9700	C21—C22	1.388 (3)
C2—C3	1.477 (3)	C22—C23	1.373 (4)
C3—C4	1.385 (3)	C22—H22	0.9300
C3—C8	1.391 (3)	C23—C24	1.363 (5)
C4—C5	1.385 (3)	C23—H23	0.9300
C4—H4	0.9300	C24—C25	1.374 (4)
C5—C6	1.387 (3)	C24—H24	0.9300
C5—H5	0.9300	C25—C26	1.380 (4)
C6—C7	1.394 (3)	C25—H25	0.9300
C6—C9	1.477 (3)	C26—H26	0.9300
C7—C8	1.361 (3)	C27—C28	1.382 (3)
C7—H7	0.9300	C27—C32	1.395 (3)
C8—H8	0.9300	C28—C29	1.385 (3)
C9—C10	1.388 (3)	C28—H28	0.9300
C9—C14	1.389 (3)	C29—C30	1.368 (4)
C10—C11	1.378 (4)	C29—H29	0.9300
C10—H10	0.9300	C30—C31	1.371 (4)
C11—C12	1.374 (4)	C30—H30	0.9300
C11—H11	0.9300	C31—C32	1.376 (3)
C12—C13	1.365 (4)	C31—H31	0.9300
C12—H12	0.9300	C32—H32	0.9300
C13—C14	1.382 (4)		
C21—P1—C15	111.69 (11)	C16—C15—P1	119.27 (19)
C21—P1—C27	106.52 (10)	C17—C16—C15	120.0 (3)
C15—P1—C27	108.17 (11)	C17—C16—H16	120.0
C21—P1—C1	109.71 (11)	C15—C16—H16	120.0
C15—P1—C1	111.96 (11)	C18—C17—C16	119.5 (3)
C27—P1—C1	108.60 (11)	C18—C17—H17	120.3
O3—S1—O2	113.5 (2)	C16—C17—H17	120.3
O3—S1—O4	113.63 (16)	C19—C18—C17	121.2 (3)
O2—S1—O4	116.7 (2)	C19—C18—H18	119.4
O3—S1—C33	104.93 (19)	C17—C18—H18	119.4

O2—S1—C33	102.5 (2)	C18—C19—C20	119.9 (3)
O4—S1—C33	103.55 (16)	C18—C19—H19	120.0
C2—C1—P1	111.90 (16)	C20—C19—H19	120.0
C2—C1—H1A	109.2	C19—C20—C15	120.0 (3)
P1—C1—H1A	109.2	C19—C20—H20	120.0
C2—C1—H1B	109.2	C15—C20—H20	120.0
P1—C1—H1B	109.2	C26—C21—C22	119.7 (2)
H1A—C1—H1B	107.9	C26—C21—P1	118.85 (18)
O1—C2—C3	122.0 (2)	C22—C21—P1	121.1 (2)
O1—C2—C1	119.1 (2)	C23—C22—C21	119.4 (3)
C3—C2—C1	118.8 (2)	C23—C22—H22	120.3
C4—C3—C8	118.4 (2)	C21—C22—H22	120.3
C4—C3—C2	123.7 (2)	C24—C23—C22	121.0 (3)
C8—C3—C2	117.9 (2)	C24—C23—H23	119.5
C5—C4—C3	120.6 (2)	C22—C23—H23	119.5
C5—C4—H4	119.7	C23—C24—C25	120.0 (3)
C3—C4—H4	119.7	C23—C24—H24	120.0
C4—C5—C6	120.9 (2)	C25—C24—H24	120.0
C4—C5—H5	119.6	C24—C25—C26	120.2 (3)
C6—C5—H5	119.6	C24—C25—H25	119.9
C5—C6—C7	117.7 (2)	C26—C25—H25	119.9
C5—C6—C9	122.2 (2)	C25—C26—C21	119.7 (2)
C7—C6—C9	120.1 (2)	C25—C26—H26	120.2
C8—C7—C6	121.5 (2)	C21—C26—H26	120.2
C8—C7—H7	119.2	C28—C27—C32	119.7 (2)
C6—C7—H7	119.2	C28—C27—P1	121.99 (18)
C7—C8—C3	120.8 (2)	C32—C27—P1	117.89 (17)
C7—C8—H8	119.6	C27—C28—C29	119.7 (2)
C3—C8—H8	119.6	C27—C28—H28	120.1
C10—C9—C14	117.9 (2)	C29—C28—H28	120.1
C10—C9—C6	122.2 (2)	C30—C29—C28	120.3 (2)
C14—C9—C6	119.9 (2)	C30—C29—H29	119.9
C11—C10—C9	120.7 (3)	C28—C29—H29	119.9
C11—C10—H10	119.7	C29—C30—C31	120.2 (2)
C9—C10—H10	119.7	C29—C30—H30	119.9
C12—C11—C10	120.3 (3)	C31—C30—H30	119.9
C12—C11—H11	119.8	C30—C31—C32	120.6 (2)
C10—C11—H11	119.8	C30—C31—H31	119.7
C13—C12—C11	119.9 (3)	C32—C31—H31	119.7
C13—C12—H12	120.1	C31—C32—C27	119.5 (2)
C11—C12—H12	120.1	C31—C32—H32	120.3
C12—C13—C14	120.1 (3)	C27—C32—H32	120.3
C12—C13—H13	119.9	F1—C33—F2	109.2 (4)
C14—C13—H13	119.9	F1—C33—F3	105.9 (4)
C13—C14—C9	120.9 (2)	F2—C33—F3	104.8 (4)
C13—C14—H14	119.5	F1—C33—S1	114.5 (3)
C9—C14—H14	119.5	F2—C33—S1	112.4 (3)
C20—C15—C16	119.4 (2)	F3—C33—S1	109.3 (3)
C20—C15—P1	121.34 (19)		

supplementary materials


Hydrogen-bond geometry (\AA , $^\circ$)

Cg1, *Cg2* and *Cg3* are the centroids of the C1–C6, C21–C26 and C15–C20 phenyl rings, respectively.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C1—H1B \cdots O4 ⁱ	0.97	2.22	3.191 (4)	178
C26—H26 \cdots O3 ⁱⁱ	0.93	2.45	3.373 (3)	174
C32—H32 \cdots O3 ⁱⁱ	0.93	2.46	3.369 (4)	168
C1—H1A \cdots Cg1 ⁱⁱⁱ	0.97	2.84	3.780 (3)	164
C10—H10 \cdots Cg2 ^{iv}	0.93	3.02	3.767 (4)	138
C23—H23 \cdots Cg3 ^v	0.93	2.91	3.788 (4)	159

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, y-1/2, -z+1/2$; (iii) $-x, -y+1, -z$; (iv) $-x, y+1/2, -z+1/2$; (v) $x, -y+1/2, z+1/2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

