

CABI is a trading name of CAB International

CABI
Nosworthy Way
Wallingford
Oxfordshire OX10 8DE
UK

CABI
745 Atlantic Avenue
8th Floor
Boston, MA 02111
USA

Tel: +44 (0)1491 832111
Fax: +44 (0)1491 833508
E-mail: info@cabi.org
Website: www.cabi.org

Tel: +1 (617) 682 9015
E-mail: cabi-nao@cabi.org

© CAB International 2016. All rights reserved. No part of this publication may be reproduced in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without the prior permission of the copyright owners.

A catalogue record for this book is available from the British Library, London, UK.

Library of Congress Cataloging-in-Publication Data

Names: Husaini, Amjad M., editor. | Neri, Davide, editor.

Title: Strawberry : growth, development and diseases / [edited by] Dr. Amjad M. Husaini and Davide Neri.

Description: Boston, MA : CABI, [2016] | Includes bibliographical references and index.


Identifiers: LCCN 2016022767 (print) | LCCN 2016023629 (ebook) | ISBN 9781780646633 (hbk : alk. paper) | ISBN 9781780646640 (ePDF) | ISBN 9781780646657 (ePub)

Subjects: LCSH: Strawberries.

Classification: LCC SB385 .S743 2016 (print) | LCC SB385 (ebook) | DDC 634/.75--dc23

LC record available at <https://lcn.loc.gov/2016022767>

ISBN-13: 978 1 78064 663 3



g. 2.1. World distribution of *Fragaria* spp.

Fragaria chiloensis (میوه درشت)



Fragaria vesca L.
قدیمی ترین (اوایل 1300)



Fragaria moschata

عطر و بوی شبیه به انگور موسکادین



Fragaria virginiana

مقاوم به سرما و خشکی



Frezier




نام علمی؟



Fragaria × ananassa Duch.





از 34 گونه، تنها دو گونه (musky , woodland or alpine) در سطح کم کشت و کار می شوند



Strawberries

- Types of strawberries
 - Junebearing
 - Day neutral
 - Alpine
 - Ornamental






Strawberries

- Self fruitful
- Cultivars
 - Junebearing
 - Early season: 'Earliglow', 'Annapolis'
 - Midseason: 'Honeoye', 'Allstar', 'Redchief', 'Guardian', 'Darselect'
 - Late season: 'Lateglow', 'Winona'
 - Day Neutral: 'Albion', 'Seascape', 'San Andreas'
 - Alpine: 'Alexandria', 'Alpine Yellow', 'Reugen'
 - Ornamental: 'Pink Panda'





8x

4X

?

?

5X



10X

2x

X

F. vesca




Fig. 1.1. Trend in strawberry area harvested across major regions. K, thousand.




Fig. 1.3. Trend in strawberry production across major regions. M, million.



NO.1 Green Strawberry 50 seeds



NO.2 Black Strawberry 50 seeds



NO.3 Red Climbing Strawberry 50 seeds



NO.4 Pineberry 80 seeds



NO.5 White Strawberry 40 seeds



NO.6 Normal Red Strawberry 50 seeds




NO.7 Bonsai Red Strawberry 50 seeds



NO.8 Cream Red Strawberry 50 seeds



NO.9 Perennial Red Strawberry 40 seeds



No.10 Giant Strawberry 50 Seeds



NO.11 Mini Red Strawberry 50 seeds



NO.12 African Blue Strawberry 50 seeds



Fig. 4.1. (a) White strawberry culture in Contulmo, Región del Bio-Bio, Chile. (b) Infected plant. (c) Commercialization of white and red strawberry at the Putu market, Región del Maule, Chile. (a, b) From Rudi Montenegro (Universidad Austral de Chile, Chile); (c) from Cristina Theoduloz (Universidad de Talca, Chile).


2013

China, mainland	2997504
United States of America	1382096
Mexico	379464
Turkey	372498
Spain	312466
Egypt	262432
Republic of Korea	216803
Poland	192647
Russian Federation	188000
Japan	165600
23. Iran (Islamic Republic of)	39296

	A	B	C	D	E
1	Area	Item	Year	Unit	Value
2	China, ma	Strawberr	2020	tonnes	3326816
3	United Sta	Strawberr	2020	tonnes	1055963
4	Egypt	Strawberr	2020	tonnes	597029
5	Mexico	Strawberr	2020	tonnes	557514
6	Turkey	Strawberr	2020	tonnes	546525
7	Spain	Strawberr	2020	tonnes	272550
8	Brazil	Strawberr	2020	tonnes	218881
9	Russian Fe	Strawberr	2020	tonnes	218400
10	Poland	Strawberr	2020	tonnes	167300
11	Morocco	Strawberr	2020	tonnes	166955
12	Japan	Strawberr	2020	tonnes	163735
13	Republic o	Strawberr	2020	tonnes	163646
14	Germany	Strawberr	2020	tonnes	152180
15	United Kir	Strawberr	2020	tonnes	129633
16	Italy	Strawberr	2020	tonnes	121790
17	Greece	Strawberr	2020	tonnes	84220
18	Netherlan	Strawberr	2020	tonnes	77570
19	Belarus	Strawberr	2020	tonnes	77059
20	Iran (Islan	Strawberr	2020	tonnes	64969


Production share of Strawberries by region

2017



Production of Strawberries: top 10 producers

2017



22	QC	Crops	102	Iran (Islam	5510	Productio	544	Strawberr	2017	2017	tonnes	55621
----	----	-------	-----	-------------	------	-----------	-----	-----------	------	------	--------	-------


Table 1.2. Nutritional composition of strawberry (*Fragaria × ananassa* Duch.). (From US Department of Agriculture: <http://ndb.nal.usda.gov/ndb/search/list?qlookup=09316&format=Full>.)

Component	Per 100 g	Standard error	Component	Per 100 g	Standard error
Nutrient			Lipids		
Water (g)	90.95	0.214	Fatty acids, total saturated (g)	0.015	–
Energy (kcal)	32	–	16:00 (g)	0.012	–
Energy (kJ)	136	–	18:00 (g)	0.003	–
Protein (g)	0.67	0.026	Fatty acids, total monounsaturated (g)	0.043	–
Total lipid (fat) (g)	0.3	0.047	16:1 undifferentiated (g)	0.001	–
Ash (g)	0.4	0.021	18:1 undifferentiated (g)	0.042	–
Carbohydrate, by difference (g)	7.68	–	Fatty acids, total polyunsaturated (g)	0.155	–
Fibre, total dietary (g)	2	0.152	18:2 undifferentiated (g)	0.09	–
Sugars, total (g)	4.89	–	18:3 undifferentiated (g)	0.065	–
Sucrose (g)	0.47	0.328	18:4 (g)	0	–
Glucose (dextrose) (g)	1.99	0.194	20:4 undifferentiated (g)	0	–
Fructose (g)	2.44	0.198	20:5 <i>n</i> -3 (EPA) (g)	0	–
Lactose (g)	0	0	22:5 <i>n</i> -3 (DPA) (g)	0	–
Maltose (g)	0	0	22:6 <i>n</i> -3 (DHA) (g)	0	–
Galactose (g)	0	0	Cholesterol (mg)	0	–
Starch (g)	0.04	0.029	Phytosterols (mg)	12	–
Vitamins			Amino acids		
<u>Vitamin C, total ascorbic acid (mg)</u>	58.8	2.473	Tryptophan (g)	0.008	–
Thiamin (mg)	0.024	0.003	Threonine (g)	0.02	–
Riboflavin (mg)	0.022	0.008	Isoleucine (g)	0.016	–
Niacin (mg)	0.386	0.037	Leucine (g)	0.034	–
Pantothenic acid (mg)	0.125	0.003	Lysine (g)	0.026	–
Vitamin B-6 (mg)	0.047	0.012	Methionine (g)	0.002	–
<u>Folate, total (µg)</u>	24	5.465	Cystine (g)	0.006	–
Folic acid (µg)	0	–	Phenylalanine (g)	0.019	–
Folate, food (µg)	24	5.465	Tyrosine (g)	0.022	–
Folate, DFE (µg)	24	–	Valine (g)	0.019	–
Choline, total (mg)	5.7	–	Arginine (g)	0.028	–
Betaine (mg)	0.2	–	Histidine (g)	0.012	–
Vitamin B-12 (µg)	0	–	Alanine (g)	0.033	–
Vitamin B-12, added (µg)	0	–	Aspartic acid (g)	0.149	–
Vitamin A, RAE (µg)	1	0.031	Glutamic acid (g)	0.098	–

Retinol (µg)	0	–	Glycine (g)	0.026	–
Carotene, β (µg)	7	0.22	Proline (g)	0.02	–
Carotene, α (µg)	0	0	Serine (g)	0.025	–
Cryptoxanthin, β (µg)	0	0	Minerals		
Vitamin A, (IU)	12	0.625	Calcium (Ca) (mg)	16	0.562
Lycopene (µg)	0	0	Iron (Fe) (mg)	0.41	0.026
Lutein + zeaxanthin (µg)	26	8.04	Magnesium (Mg) (mg)	13	0.222
Vitamin E (α-tocopherol) (µg)	0.29	0.024	Phosphorus (P) (mg)	24	0.72
Vitamin E, added (mg)	0	–	Potassium (K) (mg)	153	4.073
Tocopherol, β (mg)	0.01	0.002	Sodium (Na) (mg)	1	0.1
Tocopherol, γ (mg)	0.08	0.01	Zinc (Zn) (mg)	0.14	0.013
Tocopherol, δ (mg)	0.01	0.005	Copper (Cu) (mg)	0.048	0.004
Vitamin D (D2 + D3) (µg)	0	–	Manganese (Mn) (mg)	0.386	0.018
Vitamin D (IU)	0	–	Selenium (Se) (µg)	0.4	–
Vitamin K (phylloquinone) (µg)	2.2	0.29	Fluoride (F) (µg)	4.4	0.4
Anthocyanidins			Flavonols		
Petunidin (mg)	0.1	0.1	Isorhamnetin (mg)	0	–
Delphinidin (mg)	0.3	0.28	Kaempferol (mg)	0.5	0.01
Malvidin (mg)	0	0.01	Myricetin (mg)	0	0.04
Pelargonidin (mg)	24.8	0.69	Quercetin (mg)	1.1	0.04
Peonidin (mg)	0	0.05	Isoflavones		
Cyanidin (mg)	1.7	0.05	Daidzein (mg)	0	0
Flavan-3-ols			Genistein (mg)	0	0
(+)-Catechin (mg)	3.1	0.19	Glycitein (mg)	0	–
(–)-Epigallocatechin (mg)	0.8	0.35	Total isoflavones (mg)	0	0.005
(–)-Epicatechin (mg)	0.4	0.13	Formononetin (mg)	0	–
(–)-Epicatechin 3-gallate (mg)	0.2	0.02	Coumestrol (mg)	0	–
(–)-Epigallocatechin 3-gallate (mg)	0.1	0.06	Proanthocyanidin		
(+)-Galocatechin (mg)	0	0.005	Proanthocyanidin monomers (mg)	3.7	0.8
Flavanones			Proanthocyanidin dimers (mg)	5.3	1.89
Hesperetin (mg)	0	0	Proanthocyanidin trimers (mg)	4.9	2.27
Naringenin (mg)	0.2	0.25	Proanthocyanidin 4–6mers (mg)	28.1	6.47
Flavones			Proanthocyanidin 7–10mers (mg)	23.9	3.47
Apigenin (mg)	0	0	Proanthocyanidin polymers (>10mers) (mg)	75.8	13.36
Luteolin (mg)	0	0.001			

Functional Food

Strawberries Varieties



Junebearing

strawberries produce fruit from early June through late July and are generally overwintered

- Junebearing plants set flower buds in the fall of the year prior to fruiting when the day is relatively short, about 12 hours



Photoperiod: < 14 h or temp. < 15 C
Most sensitive to temp.

Strawberries Varieties

→ **Everbearing** strawberries produce two to three harvests of fruit intermittently during the spring, summer and fall and do not send out many runners.

• **Day Neutral** strawberries produce fruit throughout the growing season. These strawberries also produce few runners.


گھترین مساسیت به دما




• **Long day** long day induces flowering (high temp. is limiting)



زمانی که دما در محدوده 22/15(شب/روز) درجه سانتیگراد باشد، توت فرنگی بدون توجه به طول روز گلدهی خواهد داشت.



Perennial
herb
runner
triplet denticulated leaves
crown (compact stem in the
center)
daughter plant




Modified Stems

- A **stolon** is a horizontal stem that is fleshy or semi-woody and lies along the top of the ground.
- A runner is a type of stolon. Strawberry
- It is a specialized stem that grows on the soil surface and forms a new plant at one or more of its nodes.
- The leaves on strawberry runners are small but are located at the nodes which are easy to see.

STRAWBERRY PLANT

When planting strawberries, take care to get the planting depth just right. Don't bury the CROWN. Leaves, flowers, and fruit arise from there.




The Crown


- The crown is a short stem.
- New leaves develop with a bud in the axil of each leaf (auxiliary buds).
- The auxiliary buds will either remain undeveloped, grow and form runners or grow to form a side or branch crown.
- Environmental factors like day length and temperature influence the development of auxiliary buds.
- The buds of June bearers develop into runners during the summers long 14 to 16 hour days. Once the days grow shorter in August and September, the buds no longer form runners. Instead they form branch crowns and flower buds.
- The more branch crowns that are formed, the greater the potential for fruit production during the next year since each crown can produce one flower stalk on which fruit is borne.
- The buds of day neutrals can initiate flowers under any day length. For the Prairies this can range from mid June to October.



White
2.5 cm
50-500 pistils on a pulvinate yellow
receptacle





cyme inflorescence



www.knovin.ir
Strawberry inflorescence

The inflorescence develops a central or primary flower that opens first. This is followed by smaller secondary, tertiary and quaternary flowers opening up in sequence. The large primary flower develops into the largest berry, known as the "king" berry.

It usually takes about 30 days for flowers of June bearers to develop into fruit. This period may be reduced to 18 days with good growing conditions and high temperatures. Adverse conditions like cool weather could extend development to 40 days or more.




Each individual pistil (female flower part) must be pollinated


Each individual pistil develops into a "seed" that is actually a berry!




Self fruit full preferably bee



Cross-pollination by insects



Self & Wind pollination



VS.



The edible part of strawberries is the false fruit (accessory), resulting from the development of the receptacles, once fertilization occurred. The true fruits of strawberries are achenes, dry indehiscent fruits that remain attached to the fleshy false fruit. However, as we intend and eat strawberries, the fruit of the plant of strawberries is the edible part, characterized by a surface, covered with many yellow and brown small dots.



The Root


The strawberry plant is shallow rooted. Most of its roots are contained in the top 6 in (15 cm) of soil.

Day neutral plants have a shallower root system with most roots in the top 3 to 4 in (8 to 10 cm).




Planting Systems

Matted Row System



Hill System



The hill or plasticulture system uses the crowns to produce fruit with any runners removed to encourage the plant to focus on production.

Open Field Production Systems

	Hill	Matted Row
Winter temperatures	Mild	Cold
Planting dates	Summer or winter	Spring
Bed height	Raised	Flat
Mulch	Clear or black plastic	Straw
Irrigation	Trickle	Overhead
Production seasons	1–2	3–5
Cultivars	Everbearing	June bearing



Plastic covered hills ready for planting



Matted rows are planted at a lower density

Tunnel and Glasshouse Production

Tunnel production – utilizing polythene film supported by semi-circular steel frames to cover the raised beds – is widely used in Europe and the USA and helps to bring forward harvest by around 1 month and also to extend the season.

"High tunnels" - taller structures that are also enclosed against the elements - are usually passively heated and cooled, so operating costs can be minimal. Drip irrigation and plastic mulches are commonly used in crops planted into soil at high plant densities. In tunnels or glasshouses, strawberries are usually produced using the annual hill production system, spaced 30cm apart in staggered rows, with 2-4 rows in each bed (Figure 4).

Coir or coco peat – a waste fiber from coconut husks – is increasingly popular for use as a substrate in bags, slabs or pots. It needs to be prepared to remove fines and salts such as sodium chloride, which limit root development. Coir provides a better, more stable, growing media than peat and a higher level of production consistency – it is also less acid than peat at a typical pH of 4.5-6, compared to peat at 3.5-4.5. It requires an adapted fertigation programme with higher levels of calcium to balance the naturally high potassium levels in the coir, which depending upon product grade can vary from 1.6-2.0% on a dry weight basis. Coir can be used for up to four seasons and because of its more open structure water management around the roots is easier to control.



Plastic tunnel framework



High plastic tunnels

Look at all these tunnels!



How many tunnels can you see?

Matted row system of planting

- 2 feet between plants
- 3-4 feet between rows
- Watch runner placement
- Flower bud removal the 1st year for June bearing and first 60-80 days for ever-bearers



Plant in spring March-April-May



Runners develop during summer



Full "matted row" by fall



Planting depth - crown at soil level



Remove flowers during planting year



Removing the runners to maintain rows and decrease disease problems



Before



After

Getting strawberries ready for winter with mulch



Mulching strawberries

- Spread clean straw in late November or December
- 3 nights in the low 20's
- 4-6 inches deep -300 to 400 small straw bales per acre
- Rake into row when growth begins in spring
- Put back on for early frost protection-hard to do on a larger scale



Frost protection

- Watch for frost warnings whenever flowers are present
- The real need for irrigation
- Keep an eye on the Dew Point each night
- What do frosted flowers look like – black flower centers two days after a frost



Frost protection

30 days from bloom to harvest



Floating row covers

- earlier fruit harvest
- also an alternative to mulch



Renovating strawberries after harvest



1st - Mow off leaves



2nd - Narrow rows to
8-10 inches



3rd - Fertilize






Strawberries

- Junebearing culture, Year 1
 - Plant early in the spring
 - Remove flowers
 - Position runners to fill the row
 - Keep weeds under control
 - Fertilize in August
 - Mulch in late fall



Strawberries

- Junebearing culture, Year 1
 - Mulching materials
 - 4-6 inches of clean straw
 - Spunbonded row covers
 - Mulch after ground is cold (late November-December)





Strawberries

- Junebearing culture, Year 2 and beyond
 - Remove mulch in March-April
 - Harvest fruit in May-June
 - Renovate after harvest
 - Fertilize in August
 - Control weeds
 - Mulch in late fall

First, the farmer makes soil mounds.



The mounds are called beds.

Some farmers use a machine to do this.



The machine is called a bedformer.

Next, the beds are covered with plastic.



The farmer makes holes in the plastic and plants the strawberry



The plastic keeps the strawberries warm and helps them grow.

The strawberry plants grow small white flowers.



The flowers die and fruit grows in their place.



Can you see the strawberries?

What colour are they?

When the strawberries turn red, they are ready to be picked.



Climate

- وجود زمستانها و تابستانهای معتدل (cool temperate)
- دمای بهینه کمتر از سایر میوه ها (بهترین دما: 10 تا 26 درجه سلسیوس)
- کاهش کیفیت میوه با افزایش دما، افزایش تولید برگ و رانر
- احداث تونلهای پلاستیکی در مناطق سردسیر برای محافظت از یخبندان زمستانه
- استفاده از آبیاری بارانی در تابستان در مناطق گرمسیری برای محافظت از دمای بالا
- حداقل شش ساعت آفتاب کامل
- شرایط دمایی با ثبات
- از بادهای شدید در امان باشد
- در نیمکره شمالی بهترین مکان برای تولید زمینی است که شیب ملایمی (کمتر از 10 درصد) به سمت جنوب دارد.
- توت فرنگی های تازه کاشته شده به سرمای زودرس پاییزه یا بهاره حساس اند به ویژه زمانی که در مرحله رشد فعال رویشی (در مقایسه با حالت رکود زمستانه) باشند. بنابراین توصیه می شود تا عملیات پوشش گیاهان با مالچ به ویژه در مناطق سردسیرتر انجام گیرد. زمانی که دما به کمتر از 6 درجه سانتیگراد زیر صفر برسد، باید روی بوته های توت فرنگی را با پوششی از کاه و کلش پوشاند و پس از چند هفته مقدار بیشتری کاه و کلش روی گیاهان اضافه کرد (به اندازه یک پوشش 7 تا 14 سانتیمتری). این پوشش گیاهان (به ویژه جوانه ها و طوقه) را از آسیب سرما حفظ می کند.
- حساس به یخ بندان بهاره ...

Chilling requirement

ü نیاز سرمایی پایین

ü حدود 200 تا 300 ساعت دمای پایین بین صفر تا 7 درجه سانتیگراد

- کاهش نرخ رشد رویشی و زایشی (سطح برگ کوچکتر، تعداد روندکهای کمتر، کاهش تعداد گلها و گل آذینها، نرخ رشد پایین ریشه ها) در زمان رکود زمستانه
- از بین رفته طوقه در دمای 6- تا 12-
- میتوان گیاهان دختری که نیاز سرمایی آنها برطرف شده است را از مناطق سردسیرتر (مثال کردستان) تهیه کرد و سپس در مناطق گرمتر (استانهای جنوبی) کشت نمود.

Soil

- Sandy loam- sand- sandy clay
- Well drainage
- pH 5.5-6.5 (pH 5-7)

depth: Not important (short life cycle-
shallow roots)


sensitive to waterlogging
(on 15 to 25 cm hills, not heavy soils)

sensitive to different pest and disease
(not heavy soils)

sensitive to salinity

توت فرنگی نباید در مزرعه ای که در طی چهار سال گذشته در آن توت فرنگی، تمشک، یونجه، سیب زمینی، فلفل، بادمجان، لوبیا، هویج و بامیه پرورش یافته کشت گردد مگر آنکه عملیات تدخین خاک انجام شده باشد.
در کشت هیدروپونیک محلول تغذیه ای باید pH 5/7 و هدایت الکتریکی در حدود 1/5 محدود 1/7 دسی زیمنس بر متر تنظیم گردد.

Strawberry site selection

- Full sun best
- Well-drained loam to sandy loam soil
- Not after some vegetables, other berries, or sod
- Vegetables include tomato, potato, green pepper and eggplant
- Weed free area 



Soil fumigation is necessary
(soil texture is important)
Solarization
methylbromide

(۲): ویژگی‌های بهینه خاک برای کشت توت‌فرنگی

ویژگی خاک		مقادیر
pH		۵/۵-۶/۵
EC (دسی‌زیمنس بر متر)		۱/۵
مواد آلی خاک (درصد)		۲-۳
فسفر	در دسترس (کیلوگرم در هکتار)	قابل تبادل (کیلوگرم در هکتار)
پتاسیم	۶۷-۹۰	-
منیزیم	-	۳۱۵-۳۶۰
بور	-	۲۸۰
روی	۱/۷-۲/۲۵	-
	۱۱-۱۳/۵	-

Weeds are a problem in strawberries



Irrigation

- آبیاری با آب با کیفیت
- رطوبت دائمی در منطقه ریشه (حدود 90 درصد ریشه هادر عمق 7/5 تا 15 سانتیمتری سطح خاک قرار دارند)

جدول (۳): ویژگی‌های بهینه برای آب آبیاری توت‌فرنگی

ویژگی آب	سطح بهینه
pH	۶/۵-۸/۵
هدایت الکتریکی (EC)	< ۱ دسی‌زیمنس بر متر
میزان نمک‌های محلول	< ۴۵۰ میلی‌گرم بر لیتر
نرخ جذب سدیم (SAR)	< ۳۰ میلی‌گرم بر لیتر
غلظت کلر	< ۱۳۰ میلی‌گرم بر لیتر
غلظت بور	< ۰/۷ میلی‌گرم بر لیتر
غلظت نیترات	< ۵ میلی‌گرم بر لیتر
میزان بی‌کربنات	< ۱/۵ میلی‌اکی‌والان بر لیتر

جدول (۴): غلظت عناصر مختلف در تجزیه برگی توت فرنگی

غلظت عناصر پر مصرف (%)	کمبود	بهینه	زیادی
نیتروژن	$< 1/5$	$1/9 - 1/2$	> 4
نیترات		< 0.18	
فسفر	< 0.12	$0.125 - 0.14$	> 0.15
پتاسیم	$< 1/2$	$1/6 - 2/5$	$> 3/5$
کلسیم	< 0.16	$0.17 - 1/7$	> 2
منیزیم	< 0.25	$0.3 - 0.49$	> 0.18
گوگرد	< 0.12	$0.14 - 0.16$	> 0.18
سدیم		< 0.1	
کلر		< 0.15	
غلظت عناصر کم مصرف (میلی گرم بر کیلوگرم)			
منگنز	< 40	$50 - 200$	> 250
آهن	< 30	$60 - 250$	> 250
روی	< 15	$20 - 49$	> 80
مس	< 5	$7 - 19$	> 20
بور	< 19	$30 - 64$	> 90
مولیبدن	< 0.15	> 0.15	

Some farmers grow strawberries in large plastic tunnels.



The tunnels protect the strawberries from bad weather.

When the strawberries are ready, they are picked by hand.



They have to be picked by hand because they are too soft to be picked by machine.

Can you see the pickers?

Harvest

- 28 تا 34 روز پس از بازشدن گلها (در صورت بهینه بودن تمام شرایط این زمان حتی کمتر است و به حدود 24 روز میرسد)
- فاصله زمانی بین اولین گلها تا مرحله گلدهی کامل بین 11 تا 12 روز
- برداشت میوه ها به صورت یک روز در میان
- بلافاصله پس از برداشت میوه ها باید خنک شده و در انبار سرد نزدیک به صفر درجه سانتیگراد نگهداری شوند.
- میوه نافر از گراست
- شاخص های برداشت: زمانی که رنگ میوه ها کامل شد (90 درصد سطح میوه رنگ قرمز).
اسیدیته کل در حدود 6/1 درصد . میزان مواد جامد محلول آنها بالای 5/5 . نسبت مواد جامد محلول به اسیدیته کل در حدود 9/5.



Physiological disorders

- Fruit malformation

- گرده افشانی ضعیف
- صدمه به آگن ها در مراحل اولیه رشد
- کمبود بر
- دماهای خیلی بالا یا خیلی پایین (یخ بندان بهاره)
- حشرات مکند
- صدمات مکانیکی



Tarnished plant bug, *Lygus lineolaris*, feeds on the flowers, causing the berries to deform when they grow

Light frost injury will sometimes cause deformed berries, but this is usually associated with blackened centers in at least some adjacent flowers.



- Boron is the most commonly deficient micronutrient in strawberry plantings as it is very prone to leaching




stubby roots

- Poorly pollinated berries tend to be small with a rough, dimpled, seedy appearance similar to that of boron deficiency


Poor Pollination



Well Pollinated



Pest Damage



• Heat stress


- در هوای گرم و به خصوص نور مستقیم آفتاب سلول ها آسیب می بینند
- دمای بالا (بیش از 25 درجه سلسیوس) سبب افزایش سرعت تنفس و محدودیت اندازه و کیفیت میوه می شود
- تولید میوه های نرم و تیره و مستعد پوسیدگی

نایب سایبان، سرد نمودن ناحیه ریشه و جابجایی مناسب هوا



- هدایت الکتریکی بالای محلول خاک، شوری، آبیاری کم و هر بیماری ریشه که جذب آب را محدود کند، میوه کوچک و بسیار سفت ایجاد خواهد کرد.
- وجود آب خیلی زیاد، میزان کم پتاسیم و کلسیم و دمای بسیار بالا منجر به تولید میوه های نرم می شود.

Effect of salinity on strawberry growth



Propagation

- Asexual
- Sexual (several months, low yield in first year)
 - خریداری رانرهای تازه ریشه دار شده
 - گیاهان نشایی پدیدتیار شده (سرمادهی رانرهای تازه کشت شده موسوم به (Frigo فریگو)

Recent Advances in Strawberry Plug Transplant Technology

Edward F. Durner, E. Barclay Poling, and John L. Maas

Table 1. Definitions of terms specific to the commercial strawberry industry.

Term	Definition
Runner tip	An unrooted plantlet on the tip of a stolon (runner)
Plug plant	
Moduled plant	
Tip-raised potted plant	Small containerized plants produced from runner tips, generally grown for 3–5 weeks in 50-cell plastic trays with cell depths of 5–6 cm (1.9–2.4 inches), average crown diameter 8–11 mm (0.3–0.4 inches).
Tray plants	Containerized plants produced from runner tips, typically grown over a 5-month period (late July–December) in 8-, 12-, or 16-cell plastic trays with individual cell depths of 8–9 cm (3.1–3.2 inches), average crown diameter of 15–18 mm (0.6–0.7 inches).
Waiting-bed plants	Fresh-dug nursery plants with a crown diameter of 17 mm (0.7 inches) or more.
Fresh-dug bare-root plant	A field nursery-grown strawberry plant that is dug and transplanted to the production field within several days.
Frigo plants	Dormant strawberry plants stored for several months at $-1.5\text{ }^{\circ}\text{C}$ (29.3 $^{\circ}\text{F}$).



Figure 1 Strawberry Plug Plant

Producing the runner tips

Site selection and establishment

The first step in producing strawberry plug plants is to produce the runner tips. Runners are produced from mother plants, which can be grown either in a greenhouse or field. Plants can be grown in the soil or in peat-filled grow bags placed on the soil or on benches. Runner production is favored by high temperatures and long days, hence outdoor production would be limited to the summertime. With either system (outdoor or greenhouse) care should be taken to prevent runners from coming in direct contact with soil. In the greenhouse, suspended growing systems (Figures 2 & 3) are used to prevent runners from coming in contact with soil. In the field, a combination of straw mulch between rows and plastic mulch within the row will prevent runner tips from contacting the soil. In the greenhouse, the first runner tips can be expected about 8 to 10 weeks after establishing the mother plant.

Greenhouses provide the ideal setting for runner production. Greenhouses can be managed to provide ideal day time temperatures (above 75°F =24 C) and long photoperiods (about 16 hours). With supplemental heat and light, runner production can occur year round.



Figure 2 Suspended growing system



Figure 3 Suspended growing system with hanging runners

Runner tips should be harvested when root initials (little white or brown pegs, Figure 4) are present on the runner tip. Root initials should not be longer than $\frac{1}{2}$ inch. Additionally, at least two trifoliolate leaves (first leaves that appear from the runner tip) are needed and should be between $2\frac{1}{2}$ and 4 inches in length (Figure 5). Runner tips where the oldest trifoliolate leaf is larger or smaller will have limited success in establishment. Depending on individual needs and desire for uniformity of runner tips, tips are generally harvested every 10 to 14 days.



Figure 4 Root Pegs on Strawberry Runner Tips



Figure 5 Properly sized runner tip. Note 2 fully expanded trifoliolate leaves as well as extra $\frac{1}{2}$ " of runner that is used to anchor the plant (arrow).



کردستان روز کوتاه، میوه هایی با بافت نرم تولید میکند و بیشتر برای مصرف در بازار محلی مناسب است



پاروس روز کوتاه، میوه های آن بزرگ، مخروطی با رنگ قرمز روشن متمایل به نارنجی است. نسبتا میانرس بوده و به ویژه در شرایط گلخانه عملکرد خوبی دارد



کاماروسا روز کوتاه، دارای برگهای بزرگ، میوه های خوشمزه و مخروطی شکل که در اوایل باردهی وزن آن تا 44 گرم هم میرسد. برای کشت در مناطقی با زمستانهایی با روزهای کوتاه مناسب است. برای شهرهای شمالی کشور مناسب تر از نواحی جنوبی است. نسبت به کمبود کلسیم حساس است و در مقابل سفیدک و ورتیسلیوم مقاوم میباشد



سلوا روز خنثی با بوته های قوی و پرمحصول، همیشه بار و چهارفصل دارای میوه های سفت، بسیار خوش فرم و خاص که شکل و طعم آن مورد قبول هر سلیقه ای است.



کوبینالیزا روز کوتاه، میوه های آن بسیار درشت، سفت، با رنگ قرمز کم رنگ، مخروطی شانه دار با نوک نسبتا باریک است که در زمان رسیدن میوه ها سفید رنگ باقی میماند. ماندگاری بالا و و طعمی مطلوب دارد



پاجرو روز کوتاه، نوک میوه قرمز نسبتا تیره است. میوه ها درشت و مخروطی شکل می باشند.

Cultivars	Number of flowers/plant	Number of fruits/plant	Number of leaves/plant	Yield/plant (g)	Fruit volume (cm³)	Leaf length (cm)	Leaf width (cm)
Kordestan	3.5 bc	3 bc	7.75 a	28.74 d	12.3 c	3.75 b	2.8 abc
Parose	3 cd	2.75 c	6.25 bc	36.23 cd	13.9 a	3.7 bc	2.7 cb
Marak	5 a	4.5 a	6.75 abc	54.27 a	13.57 ab	3.95 a	3.05 a
Queen	4 b	3.75 ab	7 ab	43.08 bc	11.93 c	3.55 cd	2.58 c
Selva	2.25 d	2.25 c	5.5 c	26.38 d	14.07 a	3.45 d	2.75 abc
Camarosa	4.25 ab	4 a	6.5 abc	50.05 ab	12.73 bc	3.76 ab	3 ab

Means in the same column with different letters are significantly different ($P < 0.05$).

Cultivars	TSS (%)	TA (%)	TSS/TA	Fruit firmness (N)	Vitamin C (mg/100g juice)	pH
Kordestan	7.60 c	7.98 d	0.95 c	3.05 c	4.66 d	3.55 ab
Parose	6.38 d	9.08 b	0.70 e	3.65 b	6.16 b	3.68 a
Marak	7.38 c	6.85 e	1.08 b	3.75 ab	4.60 d	3.45 b
Queen	10.00 a	8.50 c	1.18 a	3.75 ab	7.05 a	3.62 a
Selva	8.05 b	9.23 b	0.87 d	3.73 ab	6.16 b	3.58 ab
Camarosa	6.43 d	9.58 a	0.67 e	3.85 a	5.28 c	3.64 a

Means in the same column with different letters are significantly different ($P < 0.05$).



Potentilla fruticosa

Strawberry 'Pink Panda'

