

On the 1-2-3-conjecture

Akbar Davoodi* and Behnaz Omoomi†

Department of Mathematical Sciences
Isfahan University of Technology
84156-83111, Isfahan, Iran

Abstract

A k -edge-weighting of a graph G is a function $w : E(G) \rightarrow \{1, \dots, k\}$. An edge-weighting naturally induces a vertex coloring c , where for every vertex $v \in V(G)$, $c(v) = \sum_{e \sim v} w(e)$. If the induced coloring c is a proper vertex coloring, then w is called a vertex-coloring k -edge-weighting (VC k -EW). Karoński et al. (J. Combin. Theory Ser. B, 91 (2004) 151–157) conjectured that every graph admits a VC 3-EW. This conjecture is known as the 1-2-3-conjecture. In this paper, first, we study the vertex-coloring edge-weighting of the Cartesian product of graphs. We prove that if the 1-2-3-conjecture holds for two graphs G and H , then it also holds for $G \square H$. Also we prove that the Cartesian product of connected bipartite graphs admits a VC 2-EW. Moreover, we present several sufficient conditions for a graph to admit a VC 2-EW. Finally, we explore some bipartite graphs which do not admit a VC 2-EW.

Keywords: Vertex-coloring edge-weighting; 1-2-3-conjecture; Edge weighting; Vertex-coloring 2-edge-weighting.

MSC: 05C15

1 Introduction

In this paper, we consider finite and simple graphs. An r -vertex coloring c of G is a function $c : V(G) \rightarrow \{1, \dots, r\}$. The coloring c is called a *proper vertex coloring* if for every two adjacent vertices u and v , $c(u) \neq c(v)$. A graph G is r -colorable if G has a proper r -vertex coloring.

A k -edge-weighting of a graph G is a function $w : E(G) \rightarrow \{1, \dots, k\}$. An edge-weighting naturally induces a vertex coloring c , where for every vertex $v \in V(G)$, $c(v) = \sum_{e \sim v} w(e)$, where $e \sim v$ means that e is an edge incident to vertex v . If the induced

*akbar.davoodi@math.iut.ac.ir

†bomoomi@cc.iut.ac.ir

coloring c is a proper vertex coloring, then w is called a *vertex-coloring k -edge-weighting* (VC k -EW). The minimum k which G has a VC k -EW is denoted by $\mu(G)$.

Obviously for a graph G with components G_1, G_2, \dots, G_t ; $\mu(G) = \max\{\mu(G_i) : 1 \leq i \leq t\}$. Also, note that the vertex-coloring k -edge-weighting is defined for graphs without K_2 as a component. Thus, we consider connected graphs with at least three vertices.

Karoński et al. in [8] introduced the concept of vertex-coloring k -edge-weighting and they proposed the following conjecture.

Conjecture 1.1. [8] (1-2-3-conjecture) For every connected graph G with at least three vertices, $\mu(G) \leq 3$.

Addario-Berry et al. [1] showed that for every connected graph with at least three vertices, $\mu(G) \leq 30$. Then Addario-Berry et al. [2] improved the bound to 16. Later, Wang and Yu [12] improved this bound to 13. Recently, Kalkowski et al. [7] showed that for every connected graph G with at least three vertices, $\mu(G) \leq 5$.

It is proved that for every 3-colorable graph G , $\mu(G) \leq 3$ [8]. In general the similar fact for 2-colorable graphs is not true. The only known families of bipartite graphs with $\mu(G) = 3$ are theta graphs $\theta(1, 4k_2 + 1, 4k_3 + 1, \dots, 4k_r + 1)$, which are graphs obtained by identifying the end-vertices of r paths of length $4k_i + 1$ [3].

In [2] Addario-Berry et al. proved that almost all graphs admit a VC 2-EW. Moreover, in [5] it was proved that the problem of existence of a VC 2-EW for a given graph is an NP-complete problem. The problem of classifying graphs which admit a VC 2-EW is an open problem. As a more general problem to VC 2-EW, in [9] Khatirinejad et al. considered the vertex-coloring $\{a, b\}$ -edge-weighting, where a and b are arbitrary real numbers. They explore classes of graphs which admit a vertex-coloring $\{a, b\}$ -edge-weighting. Furthermore, among some sufficient conditions for a bipartite graph to admit a VC $\{a, b\}$ -EW, they proposed the following question.

Question 1.2. [9] Is it true that every bipartite graph except $\theta(1, 4k_2 + 1, 4k_3 + 1, \dots, 4k_r + 1)$ and C_{4k+2} admits a VC $\{a, b\}$ -EW, for some real numbers a and b ?

Toward this problem, Havet et al. proved that every bipartite graph with minimum degree at least 3 has a VC $\{a, b\}$ -EW, for some a and b [6].

In this paper, we establish more sufficient conditions for a graph to admit a VC 2-EW, Moreover, we construct infinite family of bipartite graphs which do not admit a VC $\{a, b\}$ -EW. Thus, the answer to the question is negative.

In the following we present some theorems which are necessary to get our main results.

Theorem 1.3. [3] *If $G \not\cong K_2$ is a connected bipartite graph satisfying one of the following conditions, then $\mu(G) \leq 2$.*

- i) $\delta(G) = 1$, where $\delta(G)$ is the minimum degree of G .*
- ii) G has a part with an even number of vertices,*

Moreover, in (ii) there exists a VC 2-EW for G such that the color of every vertex in the even part is odd and the colors of other vertices are even.

The notion $N[v]$ denotes the set $N(v) \cup \{v\} = \{u : u \in V(G), uv \in E(G)\} \cup \{v\}$.

Theorem 1.4. [10] *Let $G \not\cong K_2$ be a connected bipartite graph. If one of the following conditions holds, then $\mu(G) \leq 2$.*

- i) There exists a vertex v such that $\deg(v) \notin \{\deg(u) : u \in N(v)\}$ and $G - N[v]$ is connected.*
- ii) There exists a vertex v of degree $\delta(G)$ such that $\deg(v) \notin \{\deg(u) : u \in N(v)\}$ and $G - v$ is connected.*
- iii) G is 3-connected.*

Theorem 1.5. [3] *Let P_n , C_n and K_n , $n \geq 3$, denote the path, cycle and complete graph with n vertices, respectively. Then*

$$\mu(P_n) = \begin{cases} 1 & n = 3 \\ 2 & n \geq 4 \end{cases}, \quad \mu(C_n) = \begin{cases} 2 & n \equiv 0 \pmod{4} \\ 3 & \text{otherwise} \end{cases}, \quad \text{and} \quad \mu(K_n) = 3.$$

The structure of this paper is as follows. In Section 2, we consider the 1-2-3-conjecture for the Cartesian product of graphs. First, we prove that if the 1-2-3-conjecture holds for two graphs G and H , then it also holds for $G \square H$. Then, we show that this fact concludes the trueness of 1-2-3-conjecture for the Cartesian product of some well known families of graphs. Moreover, we prove that the Cartesian product of connected bipartite graphs admits a VC 2-EW. Then, we determine $\mu(G)$ for some well known families of graphs. In Section 3, we present several sufficient conditions for a graph to admit a VC 2-EW. In section 4, we answer Question 1.2 in the negative by showing a new class of graphs admitting no VC 2-EW.

2 Cartesian product of graphs

The *Cartesian product* of two graphs G and H , written $G \square H$, is the graph with vertex set $V(G) \times V(H)$ specified by putting (u, v) adjacent to (u', v') if and only if $u = u'$ and $vv' \in E(H)$, or $v = v'$ and $uu' \in E(G)$.

The vertex-coloring $\{a, b\}$ -edge-weighting of Cartesian product of graphs are studied in [9] and through several theorems, the following result is obtained.

Theorem 2.1. [9] *If G and H are regular bipartite graphs, then the following graphs admit a VC 2-EW.*

- $K_n \square H$, if $n \geq 4$;
- $C_n \square H$, if $n \geq 4$ and $n \neq 5$;
- $G \square H$.

In this section, we prove that if the 1-2-3-conjecture holds for two graphs G and H , then it also holds for $G \square H$. Moreover, we prove that if G and H are bipartite graphs, then $\mu(G \square H) \leq 2$.

Theorem 2.2. *For every two graphs G and H , $\mu(G \square H) \leq \max\{\mu(G), \mu(H)\}$.*

Proof. Let $k = \max\{\mu(G), \mu(H)\}$ and $w_1 : E(G) \rightarrow \{1, 2, \dots, k\}$, $w_2 : E(H) \rightarrow \{1, 2, \dots, k\}$ be VC k -EW for graphs G and H , respectively. We define $w : E(G \square H) \rightarrow \{1, 2, \dots, k\}$, $w((u, v)(u', v)) = w_1(uu')$ and $w((u, v)(u, v')) = w_2(vv')$, where $u, u' \in V(G)$ and $v, v' \in V(H)$. It is easy to see that w is a VC k -EW for $G \square H$. \square

We need the following lemma to prove our main theorem in this section. In what follows $n(G)$ and $\kappa(G)$ denote the order and the vertex connectivity of G , respectively.

Lemma 2.3. [11] *For every two graphs G and H ,*

$$\kappa(G \square H) = \min\{\delta(G \square H), \kappa(H)n(G), \kappa(G)n(H)\}.$$

In the following theorem we prove that the Cartesian product of every two bipartite graphs admits a VC 2-EW.

Theorem 2.4. *If G and H are two connected bipartite graphs and $G \square H \neq K_2$, then $\mu(G \square H) \leq 2$.*

Proof. It can be seen that $G \square H$ is a connected bipartite graph if and only if G and H are connected bipartite graphs. If $n(G) = n(H) = 2$, then $G \square H \cong C_4$ and by Theorem 1.5, $\mu(G \square H) = 2$. If $n(G), n(H) > 2$ and both have a leaf vertex, then by Theorems 1.3(i) and 2.2, $\mu(G \square H) \leq 2$, otherwise $\delta(G) + \delta(H) \geq 3$. Hence, by Lemma 2.3, $\kappa(G \square H) \geq 3$ and by Theorem 1.4(iii), $\mu(G \square H) \leq 2$.

Now, let $n = n(G) > 2$ and $n(H) = 2$. Thus, $G \square H \cong G \square K_2$. Let $G^{(1)}$ and $G^{(2)}$ be the induced subgraphs representing the first and the second copy of G , respectively. To give a VC 2-EW for $G \square K_2$, first we assign weight 1 to all the edges in $G^{(1)}$ and weight 2 to all the edges in $G^{(2)}$. We denote the unweighted edge e incident to vertex $u \in V(G^{(1)})$ by e_u . Thus, for every two adjacent vertices u and v , where $u \in G^{(1)}$ and $v \in G^{(2)}$, independently to the weight of e_u , we have $c(u) \neq c(v)$. Now we assign a proper weight to the unweighted edges so that for every $uv \in E(G^{(1)}) \cup E(G^{(2)})$, $c(u) \neq c(v)$. We do this as follows.

Suppose (V_1, V_2) is the bipartition of $V(G)$ and define

$$w(e_u) = \begin{cases} 1, & \text{if } u \in V_1 \text{ with } \deg(u) \text{ even or } u \in V_2 \text{ with } \deg(u) \text{ odd;} \\ 2, & \text{if } u \in V_1 \text{ with } \deg(u) \text{ odd or } u \in V_2 \text{ with } \deg(u) \text{ even.} \end{cases}$$

Then, for vertex u in the first copy, $\sum_{e \sim u} w(e)$ is odd for $u \in V_1$ and is even for $u \in V_2$. Also, for vertex u in the second copy, $\sum_{e \sim u} w(e)$ is 0 or 1 (mod 4) for $u \in V_1$ and is 2 or 3 (mod 4) for $u \in V_2$. \square

Obviously, for every nontrivial graph G , $\mu(G) = 1$ if and only if G has no adjacent vertices with a same degree.

Proposition 2.5. *For every two graphs G and H , $\mu(G \square H) = 1$ if and only if $\mu(G) = 1$ and $\mu(H) = 1$.*

Proof. By Theorem 2.2, the condition is sufficient. Conversely, if $\mu(G \square H) = 1$, then $d_{G \square H}(u, v) \neq d_{G \square H}(u', v)$, where $uu' \in E(G)$. On the other hand $d_{G \square H}(u, v) = \deg_G(u) + \deg_H(v)$. Therefore, for every edge $uu' \in E(G)$, $\deg_G(u) \neq \deg_G(u')$. This implies $\mu(G) = 1$. Similarly, $\mu(H) = 1$. \square

3 Graphs with $\mu(G) \leq 2$

In this section, following to investigating the properties of graphs which admit a VC 2-EW, we present several sufficient conditions for a graph to admit a VC 2-EW. In the following, we consider separable graphs.

Theorem 3.1. *Let G be a graph which is the union of two simple connected graphs G_1 and G_2 such that $V(G_1) \cap V(G_2) = \{v\}$. If for all $u \in N(v)$, $2 \deg(u) \leq \deg(v)$ and for $i = 1, 2$ G_i is a cycle or $\mu(G_i) \leq 2$, then $\mu(G) \leq 2$.*

Proof. For each G_i which is not a cycle, there is a 2-edge-weighting such that $\sum_{e \sim u_1} w(e) \neq \sum_{e \sim u_2} w(e)$ whenever $u_1 u_2 \in E(G_i - v)$. For the case of a cycle, we can weight the edges starting from one adjacent to v by 2, 2, 1, 1 periodically, such that the last edge be incident to v . \square

Corollary 3.2. *Let G be a simple connected graph with the blocks B_1, B_2, \dots, B_r , $r \geq 2$, and cut vertices v_1, v_2, \dots, v_t . If for every j , $1 \leq j \leq t$, for all $u \in N(v_j)$ $2 \deg(u) \leq \deg(v_j)$ and for every i , $1 \leq i \leq r$ B_i is a cycle or $\mu(B_i) \leq 2$, then $\mu(G) \leq 2$.*

A *simple path* in graph G is a path P in G which the degree of each vertex of P in G is two except the two end-vertices of P . The edge set of G is disjoint union of maximal simple paths. Note that a cycle in a graph G with at most one vertex of degree greater than two in G is considered as a maximal simple path in G . In the following theorem we give several sufficient conditions for a graph to admit a VC 2-EW with respect to length of maximal simple paths in G .

Theorem 3.3. *Let G be a connected graph which is not a cycle. If one of the following conditions holds, then $\mu(G) \leq 2$.*

- i) G contains no maximal simple path of length $3 \pmod{4}$ and for every edge $e = xy$ which is a maximal simple path of G , $\deg(x) \neq \deg(y)$.*
- ii) G contains no edge as a maximal simple path.*

Proof. In each case, we show that there exist a 2-edge-weighting for maximal simple paths in G which induces a proper coloring of G .

i) Starting from an end-edge, we assign the weight 1 and 2 to the edges of maximal simple paths of length $2 \pmod{4}$, according to the pattern 2, 2, 1, 1, periodically. Also for maximal simple paths of length $0, 1 \pmod{4}$, we follow the pattern 2, 1, 1, 2.

According to the given weighting, for two arbitrary adjacent vertices u and v , clearly if the degree of one of these vertices is one, then $c(u) \neq c(v)$. Moreover, if the degrees of both vertices u and v are two, then either u and v are internal vertices of a maximal simple path and $c(u) \neq c(v)$.

Now, assume that $\deg(v) \geq 3$ that means v is end-vertex of a maximal simple path. By the given patterns and by the assumption, every edge incident to v has weight 2, thus $c(v) = 2 \deg(v) \geq 6$. If u is an internal vertex for a maximal simple path, then $c(u) \in \{3, 4\}$ and $c(u) \neq c(v)$. If $e = uv$ is a maximal simple path of G , then by assumption $c(u) = 2 \deg(u) \neq 2 \deg(v) = c(v)$.

ii) In this case, similar to above, we weight the edges of maximal simple paths of length $0, 1, 3 \pmod{4}$, by the pattern $2, 1, 1, 2$. For maximal simple paths of length $2 \pmod{4}$, we follow the pattern $1, 1, 2, 2$.

With a similar discussion as the above one for adjacent vertices u and v both of degree less than three, we have done. Thus, let $\deg(v) \geq 3$. By the assumption, for every vertex $u \in N(v)$, $\deg(u) = 2$, hence $c(u) \in \{2, 3, 4\}$. Clearly if $\deg(v) > 4$ or $c(v) > 4$, then $c(u) \neq c(v)$. If $\deg(v) = c(v) = 4$, then by the given patterns, $c(x) = 2$ for every $x \in N(v)$. Thus, particularly $c(u) \neq c(v)$. If $\deg(v) = 3$, then there are two possibilities $c(v) = 3$ and $c(v) = 4$. If $c(v) = 3$, then again by the given patterns, $c(u) = 2 \neq c(v)$. For the case $\deg(v) = 3$ and $c(v) = 4$, one of the edges incident to v belongs to a maximal simple path, say p_0 , has weight 2 and two other edges incident to v , have weight 1. Now, $c(u) \neq c(v)$ unless $u \in p_0$ and p_0 is of length $1 \pmod{4}$. Note that the two other paths incident to v , say p_1 and p_2 , are of length 2 or $3 \pmod{4}$.

If p_0 is of length $1 \pmod{4}$ and $c(u) = c(v)$ and one of the paths p_1 or p_2 , say p_1 , is of length $3 \pmod{4}$, then by reweighting the edges of p_1 by the same pattern but starting from v , we get $c(v) = 5 > c(x)$ for every vertex $x \in N(v)$. In this process no new adjacent vertices with a same color occurs.

If $u \in p_0$, p_0 is of length $1 \pmod{4}$, $c(u) = c(v)$ and both p_1 and p_2 are of length $2 \pmod{4}$, then we reweight the edges of p_0 by the same pattern as before but starting from v , thus we get $c(v) \neq c(x)$ for every vertex $x \in N(v)$. We have no new adjacent vertices with the same color, unless the other end-vertex of p_0 , say v' , has the conditions exactly the same as v . That means $\deg(v') = 3$ and $c(v') = 4$. By the same argument, we solve the problem when at least one of the paths incident to v' is of length $3 \pmod{4}$.

Otherwise, both of the other paths incident to v' are of length $2 \pmod{4}$. In this case we choose one of them and change its edge weight pattern from $1, 1, 2, 2$ to $2, 2, 1, 1$. If the other end of this path, say v'' , is not of color 4 or is not of degree 3, we have done as above. Otherwise we do this process for v'' .

Since the graph is finite, after iterating this process we face to an end-vertex of a maximal simple path for the second time. While the color of this vertex, after changing the weights according to the given pattern in the second visit is at least five. Thus, this process terminates and in this manner, we reduce the number of adjacent vertices with the same color to get the desired weighting. \square

For the case that G contains an edge as a maximal simple path there are graphs with $\mu(G) = 2$ and $\mu(G) = 3$. The complete graph K_n is an example of graphs in which every maximal simple path is of length 1 and $\mu(K_n) = 3$. The following proposition gives an example of graphs in which every maximal simple path is of length 1 and $\mu(G) = 2$.

Proposition 3.4. *For every complete r -partite graph $K_{r \times n}$, $r, n \geq 2$, $\mu(K_{r \times n}) = 2$.*

Proof. Let A be a weighted adjacency matrix of $G = K_{r*n}$, defined as follows.

$$A = \begin{pmatrix} 0 & B & \cdots & B \\ B^T & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & B \\ B^T & \cdots & B^T & 0 \end{pmatrix}_{rn \times rn}, \text{ where } B = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \vdots & \vdots & & \vdots \\ 1 & 1 & \cdots & 1 \\ 2 & 2 & \cdots & 2 \end{pmatrix}_{n \times n}$$

Note that the induced color on every vertex is the sum of entries in it's corresponding row in A . Thus, for every two arbitrary vertices u and v in i -th part and j -th part of G , $c(u) \in \{(r-i)n + (i-1)(n+1), 2(r-i)n + (i-1)(n+1)\}$ and $c(v) \in \{(r-j)n + (j-1)(n+1), 2(r-j)n + (j-1)(n+1)\}$. Since $i \neq j$ we have

$$(r-i)n + (i-1)(n+1) \neq (r-j)n + (j-1)(n+1)$$

and

$$2(r-i)n + (i-1)(n+1) \neq 2(r-j)n + (j-1)(n+1)$$

On the other hand, $j \leq r$ hence, $j-i \leq n(r-i)$ with equality if and only if $n = 1$ and $j = r$. Thus $(r-j)n + (j-1)(n+1) \neq 2(r-i)n + (i-1)(n+1)$. Similarly $(r-i)n + (i-1)(n+1) \neq 2(r-j)n + (j-1)(n+1)$. Therefore, A is a desired edge-weighting. \square

We end this section with the following conjecture.

Conjecture 3.5. Let $G \cong C_{4k+r}$, $r \in \{1, 2, 3\}$, be a connected graph. If $\mu(G) > 2$, then G has two adjacent vertices of a same degree greater than two.

4 Bipartite graphs

It is known that every 3-colorable graph admits a VC 3-EW [8], and there are bipartite graph G with $\mu(G) = 2$ and also bipartite graph G with $\mu(G) = 3$. The only known bipartite graphs with $\mu(G) = 3$ are C_{4k+2} and the theta graphs $\theta(1, 4k_2 + 1, \dots, 4k_r + 1)$ [3]. The *theta graph* $\theta(l_1, l_2, \dots, l_r)$ is the graph obtains from r disjoint paths, of lengths l_1, l_2, \dots, l_r , respectively, by identifying their end-vertices called *the roots* of the graph. Notice that $\theta(l_1) = P_{1+l_1}$ and $\theta(l_1, l_2) = C_{l_1+l_2}$.

Theorem 4.1. [3] *Let $G = \theta(l_1, l_2, \dots, l_r)$, where $r \geq 3$ and $1 \leq l_1 \leq l_2 \leq \dots \leq l_r$. Then,*

$$\mu(G) = \begin{cases} 1 & l_i = 2 \text{ for all } i \\ 3 & l_1 = 1 \text{ and } l_i \equiv 1 \pmod{4} \text{ for all } i \neq 1 \\ 2 & \text{otherwise.} \end{cases}$$

In this section, we consider a generalization of theta graphs, called generalized polygon trees, and answer to Question 1.2 in negative.

The class of *generalized polygon trees* is defined recursively as follows. A cycle C_p ($p \geq 3$) is a generalized polygon tree. Next, suppose H is a generalized polygon tree

containing a simple path P_k , where $k \geq 1$. The graph obtained from the disjoint union of H and a cycle C_r , where $r > k$, by identifying P_k in H with a path of length k in C_r , is also a generalized polygon tree.[4]

In Theorem 4.1, $\mu(G)$ is determined whenever G is a generalized polygon tree with two interior regions.

In this section, we determine the $\mu(G)$ for bipartite generalized polygon trees with three interior regions (see Figure 1).

We need the following sufficient condition to guarantee that a bipartite graph admits a VC 2-EW.

Theorem 4.2. *Let G be a connected bipartite graph. If G contains a vertex v such that $\deg(v) > \deg(u)$ for every $u \in N(v)$ and $G - v$ is connected, then $\mu(G) \leq 2$.*

Proof. Let U and W be the parts of G . If either $|U|$ or $|W|$ is even, then by Theorem 1.3(ii), the result follows. Thus, we assume that both parts of G are of odd size. Let $v \in U$ satisfy $\deg(v) > \deg(u)$ for every $u \in N(v)$ and $G - v$ is connected. Since $|U - v|$ is even, by Theorem 1.3(ii), $G - v$ admits a VC 2-EW such that $c(x)$ is odd for every $x \in U - v$ and $c(y)$ is even for every $y \in W$. Now we assign weight 2 to all the edges that are adjacent to vertex v . Clearly $c(x)$ is odd for every $x \in U - v$ and $c(y)$ is even for every $y \in W$. Also $c(v) = 2 \deg_G(v) > 2 \deg_G(u) \geq c(u)$ for every $u \in N(v)$. \square

Fact 4.3. *In every VC 2-EW of a path, the weight of one edge forces the weights of every second edge. Thus, if P is a path of length l , $l \equiv 1 \pmod{4}$, then two end-edges of P get a same weight and if $l \equiv 3 \pmod{4}$, then they get different weights. Furthermore, when $l \not\equiv 1, 3 \pmod{4}$ every arbitrary assignment of the end-edges of P with $\{1, 2\}$, could be extended to a VC 2-EW for P .*

Theorem 4.4. *Let G be a bipartite generalized polygon tree with at most three interior regions shown in Figure 1, where p_1, p_2, \dots, p_6 are paths of length a, b, c, d, e and f , respectively. Then $\mu(G) = 3$ if and only if one of the following conditions holds.*

- $a = b = c = d = 0$ and $e + f \equiv 2 \pmod{4}$.
- $b = 1, c = d = 0$ and $a, e + f \equiv 1 \pmod{4}$.
- $b = 1, e = f = 0$ and $a, c, d \equiv 1 \pmod{4}$.
- $b = c = 1, a, d, e \equiv 1 \pmod{4}$ and $f \equiv 3 \pmod{4}$.

Proof. We know that a graph admits a vertex-coloring 1-edge-weighting if and only if every two adjacent vertices have different degrees. It can be verified that this condition holds if and only if $a = b = c = d = e = f = 2$ or $a = d = f = 2, b = c = 1$ and $e = 0$ or $a = d = f = 2, b = c = 2$ and $e = 0$ or $a = b = c = d = 2$ and $e = f = 0$.

For b and c there are three possibilities $b = c = 0$ or $b \neq 0, c = 0$ or $b, c \neq 0$.

Let $b = c = 0$. If $a = d = 0$, then G is an even cycle C_{e+f} and by Theorem 1.5, if $e + f \equiv 0 \pmod{4}$, then $\mu(G) = 2$, otherwise, $\mu(G) = 3$. If $a \neq 0$ or $d \neq 0$, by Corollary 3.2, $\mu(G) = 2$.




Figure 1: A generalized polygon tree with three interior regions.

Now let $b \neq 0$ and $c = 0$. Since $d \neq 0$, G is a graph with two blocks: a $\theta(a, b, e + f)$ and a cycle C_d . Thus by Theorem 3.1, $\mu(G) = 2$ unless $\mu(\theta(a, b, e + f)) = 3$. In this case we give a VC 2-EW for G in Figure 2(i). In the figures the edges and paths are denoted by straight lines and curves, respectively, and every path is weighted periodically by the given pattern through the denoted direction.

Otherwise $b, c \neq 0$. In this case, we consider three possibilities $b, c > 1$ or $b = 1, c > 1$ or $b = c = 1$.

Now let $b, c > 1$. In this case if $e \neq 1$ or $f \neq 1$, then there exists a vertex x with $\deg(x) \geq 3$ such that $\deg(x) > \deg(y)$ for all $y \in N(x)$ and $G - x$ is connected. Thus by Theorem 4.2, $\mu(G) \leq 2$. Therefore we assume that $e = f = 1$. Notice that if $a = 0$ or $d = 0$, then G is a theta graph and we have done. Now, since G is bipartite $b + c$ is even. Hence, $b + c \equiv 0 \pmod{4}$ or $b + c \equiv 2 \pmod{4}$. For the first possibility if there is a part of even order, by Theorem 1.3(ii), we have done. Otherwise we have two odd parts and $a + d \equiv 2 \pmod{4}$. In this case we have four possibilities $a \equiv 2 \pmod{4}$, $b, c, d \equiv 0 \pmod{4}$ or $a, b, c \equiv 2 \pmod{4}$, $d \equiv 0 \pmod{4}$ or $a, b, d \equiv 1 \pmod{4}$, $c \equiv 3 \pmod{4}$ or $a, c, d \equiv 3 \pmod{4}$, $b \equiv 1 \pmod{4}$, in which in each case the given pattern in Figure 2(ii) is a VC 2-EW for G . For the latter case, $a + d \equiv 2 \pmod{4}$ give a part with even number of vertices and if both parts have odd number of vertices and $a + d \equiv 0 \pmod{4}$ by the symmetry of b, c and a, d with the same discussion we get the desired result.

Now let $b = 1, c > 1$. If $e = f = 0$, then $G \cong \theta(a, 1, c, d)$. Thus, we assume that $\{e, f\} \neq \{0\}$. In this case if $e \neq 1$ or $f \neq 1$, then there exists a vertex x with $\deg(x) \geq 3$ such that $\deg(x) > \deg(y)$ for all $y \in N(x)$ and $G - x$ is connected. Thus by Theorem 4.2, $\mu(G) \leq 2$. Therefore, we assume $e = f = 1$. Since G is bipartite a, c and d are odd. For the cases $a, c, d \equiv 3 \pmod{4}$ and $a \equiv 3 \pmod{4}$, $c, d \equiv 1 \pmod{4}$ and $a, c \equiv 1 \pmod{4}$, $d \equiv 3 \pmod{4}$ in Figure 2(iii) is given a VC 2-EW. Otherwise, G has a part of even order and by Theorem 1.3(ii), $\mu(G) \leq 2$.

Now if $b = c = 1$, then since G is bipartite, $e + f$ is even and a, d are odd.

Case 1. $a, d \equiv 1 \pmod{4}$. If $e + f \equiv 2 \pmod{4}$, then G has a part of even number of vertices and by Theorem 1.3(ii), $\mu(G) \leq 2$. Thus, we assume $e + f \equiv 0 \pmod{4}$. For the cases $e, f \equiv 0 \pmod{4}$ and $e, f \equiv 2 \pmod{4}$ we give a VC 2-EW for G in Figures 2(iv) and 2(v), respectively.

For the case $e \equiv 1, f \equiv 3 \pmod{4}$ first we show that $\mu(G) \geq 3$. Suppose to the

contrary that G admits a VC 2-EW. Since $a \equiv 1 \pmod{4}$, by Fact 4.3, in p_1 the incidence edges on u and v have a same weight. Similarly incidence edges on u' and v' in p_4 . Thus, the incidence edges on u, v through p_5 and p_6 must have different weights. On the other hand, since $e \equiv 1 \pmod{4}$, two end-edges on p_5 get the same weight but two edges on p_6 get different weights, because $f \equiv 3 \pmod{4}$. Therefore $c(u') = c(v')$. This contradiction implies $\mu(G) \geq 3$. On the other hand, G is bipartite, thus is 3-colorable and $\mu(G) \leq 3$. Therefore, $\mu(G) = 3$.

Case 2. $a, d \equiv 3 \pmod{4}$. If $e + f \equiv 2 \pmod{4}$, then by Theorem 1.3(ii), $\mu(G) \leq 2$. Thus, we assume $e + f \equiv 0 \pmod{4}$. Now one of the three possibilities $e, f \equiv 0 \pmod{4}$ or $e, f \equiv 2 \pmod{4}$ or $e \equiv 1, f \equiv 3 \pmod{4}$ occurs, in which in each case the given pattern in Figure 2(iv) is a VC 2-EW for G .

Case 3. $a \equiv 1, d \equiv 3 \pmod{4}$. If $e + f \equiv 0 \pmod{4}$, then by Theorem 1.3(ii), $\mu(G) \leq 2$. Thus, we assume $e + f \equiv 2 \pmod{4}$ and for the cases $e, f \equiv 1 \pmod{4}$ or $e \equiv 0 \pmod{4}, f \equiv 2 \pmod{4}$ and the case $e, f \equiv 3 \pmod{4}$ we give a VC 2-EW for G in Figures 2(v) and 2(iv), respectively. Notice that in the case $e \equiv 0 \pmod{4}, f \equiv 2 \pmod{4}$ if $e = 0$, then by Theorem 4.2, $\mu(G) \leq 2$. Also in the case $e, f \equiv 1 \pmod{4}$ if $e = 1$, we can replace weight of the edge uv with 1 to get a VC 2-EW.




Figure 2: A VC 2-EW for G .

□

In Theorem 4.4 we obtained infinite classes of bipartite graphs with $\mu(G) = 3$. It can be seen that the first three cases of such graphs are C_{4k+2} , $\theta(1, 4k_2 + 1, 4k_3 + 1)$ and $\theta(1, 4k_2 + 1, 4k_3 + 1, 4k_4 + 1)$, respectively. Furthermore, The graphs satisfying the fourth condition can be generalize to the following example which provides more bipartite graphs with $\mu(G) = 3$.

Remark 4.5. Fact 4.3 is true the edges are weighted with every two arbitrary real numbers a and b . Thus, the proof of Theorem 4.4 can be applied to characterize generalized polygon trees with at most three interior regions that has no VC $\{a, b\}$ -EW.

Example 4.6. Let G be a graph shown in Figure 3, in which Q_i , $1 \leq i \leq n + 1$, are paths of length $1 \pmod{4}$ and R_i, S_i , $1 \leq i \leq n$, are paths of odd lengths r_i, s_i , where $r_i + s_i \equiv 0 \pmod{4}$. By Remark 4.5, it is easy to see that G has no VC $\{a, b\}$ -EW. Thus since G is bipartite, $\mu(G) = 3$.




Figure 3: A family of 2-connected bipartite graphs with no VC $\{a, b\}$ -EW.

Note that in graph in Figure 3, we can add any number of paths of length $1 \pmod{4}$ to G with the same end-vertices with Q_i to get more bipartite graphs with no VC $\{a, b\}$ -EW.

The structures of known bipartite graphs with no VC $\{a, b\}$ -EW, encourage us to ask the following question.

Question 4.7. Is it true that every bipartite graph except $\theta(1, 4k_2+1, 4k_3+1, \dots, 4k_r+1)$, C_{4k+2} and generalized polygon trees described in Example 4.6 admits a VC $\{a, b\}$ -EW, for some real numbers a and b ?

Acknowledgement

The authors would like to thank the referees for their valuable comments on our submitted manuscript.

References

- [1] L. Addario-Berry, K. Dalal, C. McDiarmid, B.A. Reed, A. Thomason. Vertex-coloring edge-weightings, *Combinatorica*, 27(1):1–12, 2007.
- [2] L. Addario-Berry, K. Dalal, B.A. Reed. Degree constrained subgraphs, *Discrete Appl. Math.*, 156:1168–1174, 2008.
- [3] G.J. Chang, C. Lu, J. Wu, Q. Yu. Vertex coloring edge weighting of graphs, *Taiwanese J. Math.*, 15(4):1807–1813, 2011.

- [4] F. M. Dong, K. M. Koh, K. L. Teo. *Chromatic Polynomials and Chromaticity of Graphs*, World Scientific Publishing Company, 2005.
- [5] A. Dudek, D. Wajc. On the complexity of vertex-colouring edge-weightings, *Discrete Math. Theor. Comput. Sci.*, 13(3):45–50, 2011.
- [6] F. Havet, N. Paramaguru, R. Sampathkumar. Detection number of bipartite graphs and cubic graphs, *Rapport de recherche RR-8115, INRIA*, October 2012.
- [7] M. Kalkowski, M. Karoński, F. Pfender. Vertex-coloring edge-weightings: towards the 1-2-3-conjecture, *J. Combin. Theory Ser. B*, 100:347–349, 2010.
- [8] M. Karoński, T. Łuczak, A. Thomason. Edge weights and vertex colors, *J. Combin. Theory Ser. B*, 91:151–157, 2004.
- [9] M. Khatirinejad, R. Naserasr, M. Newman, B. Seamone, B. Stevens., Vertex-colouring edge-weightings with two edge weights, *Discrete Math. Theor. Comput. Sci.*, 14(1):1–20, 2012.
- [10] H. Lu, Q. Yu, C.-Q. Zhang. Vertex-coloring 2-edge-weighting of graphs, *European J. Combin.*, 32:21–27, 2011.
- [11] S. Špacapan. Connectivity of Cartesian products of graphs, *Appl. Math. Lett.*, 21(7):682–685, 2008.
- [12] T. Wang, Q.L. Yu. On vertex-coloring 13-edge-weighting, *Front. Math. China*, 3:581–587, 2008.